

525 Rec'd PCT/PTO 01 DEC 2000

FORM PTO-1390 (REV. 1-98)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 427.038	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 09/701893	
INTERNATIONAL APPLICATION NO. PCT/FR99/01422		INTERNATIONAL FILING DATE June 15, 1999		PRIORITY DATE CLAIMED June 15, 1998	
TITLE OF INVENTION USE OF DIAZEPINES FOR THE PREPARATION OF MEDICAMENTS INTENDED TO TREAT THE PATHOLOGICAL STATES OR THE DISEASES IN WHICH ONE OF SOMATOSTATIN RECEPTORS IS INVOLVED					
APPLICANT(S) FOR DO/EO/US BIGG et al					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). 4. <input type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> has been transmitted by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 					
Items 11. to 16. below concern document(s) or information included:					
<ol style="list-style-type: none"> 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: Modified Sheets 43-57 in English; Modified Sheets 43-57 in French; PCT/IB/332 					

U.S. APPLICATION NO. (if known, see 37 CFR 1.53)

INTERNATIONAL APPLICATION NO.

ATTORNEY'S DOCKET NUMBER

09/701893

PCT/FR99/01422

427.038

17. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$1070.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO \$930.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$790.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$720.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4) \$98.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$1000.00

\$1000.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$

CLAIMS

NUMBER FILED

NUMBER EXTRA

RATE

\$

Total claims

- 20 =

x \$22.00

\$

Independent claims

- 3 =

x \$82.00

\$

MULTIPLE DEPENDENT CLAIM(S) (if applicable)

+ \$270.00

\$

TOTAL OF ABOVE CALCULATIONS =

\$1000.00

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

\$

SUBTOTAL =

\$1000.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

TOTAL NATIONAL FEE =

\$1000.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

\$

40.00

TOTAL FEES ENCLOSED =

\$1040.00

Amount to be
refunded:

\$

charged:

\$

a. ☒ A check in the amount of \$ 1040.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 02-2275 A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR
1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Bierman, Muserlian and Lucas
600 Third Avenue
New York, NY 10016



20311

PATENT TRADEMARK OFFICE

SIGNATURE

Charles A. Muserlian

NAME

19,683

REGISTRATION NUMBER

Our Ref.: 427.038

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: :
BIGG et al : PCT Date: June 15, 1999
PCT/FR99/01422 :
Serial No.: :
Filed: Concurrently Herewith :
For: USE OF...INHIBITING FACTOR :
RECEPTORS :
600 Third Avenue
New York, NY 10016

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend this application as follows:

IN THE SPECIFICATION:

Page 1, before line 1, insert

--This application is a 371 of PCT/FR99/01422 filed June
15, 1999.--

IN THE CLAIMS:

Modified claim 1, cancel line 1 and insert --A method for
blocking somatostatin receptors in warm-blooded animals comprising
administering to warm-blooded animals an amount effective for
blocking somatostatin receptors of a compound of the formula--.

Claims 2 and 3, cancel line 1 of each and insert --The method of claim 1 wherein--.

Claim 4, cancel lines 1 and 2 and insert --The method of claim 1 wherein the compound is selected from the group consisting of--.

Claim 5, cancel line 1 and insert --A compound of the formula--.


Cancel claims 7 and 8 and add the following claim.

--9. A composition for blocking somatostatin receptors comprising an amount of a compound of claim 1 sufficient to block somatostatin receptors and an inert pharmaceutical carrier.--

REMARKS

The amendment is submitted to refer to the PCT application and to conform the method of use claims to the American practice.

Respectfully submitted,
BIERMAN, MUSERLIAN AND LUCAS


Charles A. Muserlian, #19,683
Attorney for Applicant(s)
Tel. # (212) 661-8000

CAM:sd
Enclosure: Return Receipt Postcard

Use of diazepines for the preparation of medicaments intended to treat the pathological states or the diseases in which one of the somatostatin receptors is involved

The present invention relates to the use of pyrido-thieno-triazolo-diazepines for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors are involved. The invention also relates to new pyrido-thieno-triazolo-diazepines and the therapeutic compositions containing them.


5 Somatostatin (SST) is a cyclic tetradecapeptide which was isolated for the first time from the hypothalamus as a substance which inhibits the growth hormone (Brazeau P. et al., *Science* 1973, 179, 77-79). It also operates as a neurotransmitter in the brain (Reisine T. et al., *Neuroscience* 1995, 67, 777-790; Reisine et al., *Endocrinology* 1995, 16, 427-442). Molecular cloning has allowed it to be shown that the bioactivity of somatostatin depends directly on a
10 family of five receptors.

Among the pathological disorders associated with somatostatin (Moreau J.P. et al., *Life Sciences* 1987, 40, 419; Harris A.G. et al., *The European Journal of Medicine*, 1993, 2, 97-105), there can be mentioned for example: acromegalia, hypophyseal adenomas, Cushing's disease, gonadotrophinomas and prolactinomas, catabolic side-effects of glucocorticoids,
15 insulin dependent diabetes, diabetic retinopathy, diabetic nephropathy, hyperthyroidism, gigantism, endocrinic gastroenteropancreatic tumors including carcinoid syndrome, VIPoma, insulinoma, nesidioblastoma, hyperinsulinemia, glucagonoma, gastrinoma and Zollinger-Ellison's syndrome, GRFoma as well as acute bleeding of the esophageal varices, gastroesophageal reflux, gastroduodenal reflux, pancreatitis, enterocutaneous and pancreatic
20 fistulae but also diarrheas, refractory diarrheas of acquired immunodeficiency syndrome, chronic secretory diarrhea, diarrhea associated with irritable bowel syndrome, disorders linked with gastrin releasing peptide, secondary pathologies with intestinal grafts, portal hypertension as well as hemorrhages of the varices in patients with cirrhosis, gastro-intestinal hemorrhage, hemorrhage of the gastroduodenal ulcer, Crohn's disease, systemic sclerosis, dumping
25 syndrome, small intestine syndrome, hypotension, scleroderma and medullar thyroid carcinoma, illnesses linked with cell hyperproliferation such as cancers and more particularly breast cancer, prostate cancer, thyroid cancer as well as pancreatic cancer and colorectal cancer, fibroses and more particularly fibrosis of the kidney, fibrosis of the liver, fibrosis of the lung, fibrosis of the skin, also fibrosis of the central nervous system as well as that of the nose and
30 fibrosis induced by chemotherapy, and other therapeutic fields such as, for example, cephalas including cephalas associated with hypophyseal tumors, pain, panic attacks,

chemotherapy, cicatrization of wounds, renal insufficiency resulting from delayed development, obesity and delayed development linked with obesity, delayed uterine development, dysplasia of the skeleton, Noonan's syndrome, sleep apnea syndrome, Graves' disease, polycystic disease of the ovaries, pancreatic pseudocysts and ascites, leukemia, 5 meningioma, cancerous cachexia, inhibition of H pylori, psoriasis, as well as Alzheimer's disease. Osteoporosis can also be mentioned.

In the Patent FR 2645153, the Applicant described diazepines which have an anti-PAF activity. The Applicant found that these products and products related to these compounds have an affinity and a selectivity for the somatostatin receptors. As somatostatin and its 10 peptide analogues often have a poor bioavailability by oral route and a low selectivity (Robinson, C., Drugs of the Future, 1994, 19, 992; Reubi, J.C. et al., TIPS, 1995, 16, 110), said compounds, non-peptide agonists or antagonists of somatostatin, can be advantageously used to treat the pathological states or the diseases as presented above and in which one (or more) somatostatin receptors are involved. Preferably, these compounds can be used for the 15 treatment of acromegalia, hypophyseal adenomas or endocrine gastroenteropancreatic tumors including carcinoid syndrome.

Therefore a subject of the invention is the use of a compound of general formula I



in which

20 W represents the hydrogen atom or a radical of formula R-X-C(Y)-;

R represents an aryl or heteroaryl radical, the aryl and heteroaryl radicals being optionally substituted;

X represents a radical of formula $-(CH_2)_n-Z$ in which Z represents a covalent bond, NH, O or S and n an integer of 0 to 2;

25 Y represents O or S;

R₁ represents one or more groups, identical or different, chosen from: the hydrogen atom, the hydroxy, halo radical, a lower alkyl, lower alkoxy radical;

R_{2a} and R_{2b} represent, independently, the hydrogen atom; a lower alkyl, lower alkenyl or lower alkynyl radical, the alkyl, alkenyl and alkynyl radicals being optionally substituted; or an R₂₁Z₂₁- radical in which Z₂₁ represents O, C(O), OC(O), S, and R₂₁ represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical;

R₃ represents the hydrogen atom, the halo, nitro or cyano radical, an alkyl radical with 1 to 10 carbon atoms, lower alkenyl, lower alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, lower aryloxyalkyl, heteroaryl or heteroarylalkyl, the alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted; or an R₃₁Z₃₁- radical in which Z₃₁ represents O, C(O), OC(O), S, and R₃₁ represents the hydrogen atom, a lower alkyl, aryl or lower arylalkyl radical;

or a salt of this product (when the molecule contains a basic group), for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved.

A more particular subject of the invention is the use, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved, of a compound of general formula I as defined above in which

W represents the hydrogen atom or a radical of formula R-X-C(Y)-;

R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals: lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethyloxy, hydroxy, nitro, cyano, aryl, aryloxy, cycloalkyl or heterocycloalkyl;

R₁ represents one or more identical or different groups, chosen from: the hydrogen atom, the hydroxy radical, halo, a lower alkyl radical, lower alkoxy, the alkyl and alkoxy radicals being optionally substituted by one or more identical or different radicals chosen from the following radicals: trifluoromethyl, lower alkoxy, amino, lower alkyl amino and lower dialkyl amino;

R_{2a} and R_{2b} represent, independently:

the hydrogen atom;

a lower alkyl, lower alkenyl or lower alkynyl radical, the alkyl, alkenyl and alkynyl radicals being optionally substituted by one or more identical or different radicals, chosen from:

5 halo; an -NR₂₂R₂₃ radical in which R₂₂ and R₂₃ represent, independently, the hydrogen atom, a lower alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl, arylsulphonyl, lower alkoxy carbonyl, aryloxy carbonyl, alkylcarbonyl, arylcarbonyl or cycloalkylcarbonyl radical; or a -Z₂₂R₂₄ radical in which
10 Z₂₂ represents O, S, C(O), OC(O) and R₂₄ represents a hydrogen atom, a lower alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl or arylsulphonyl radical;

an R₂₁Z₂₁- radical in which Z₂₁ represents O, C(O), OC(O), S, and R₂₁ represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical;

15 R₃ represents:

the hydrogen atom, the halo, nitro or cyano radical;

an alkyl radical with 1 to 10 carbon atoms, lower alkenyl, lower alkynyl, cycloalkyl, cycloalkylalkyl, aryl, lower arylalkyl, lower aryloxyalkyl, heteroaryl or lower heteroarylalkyl, the alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl
20 radicals being optionally substituted by one or more identical or different radicals chosen from:

halo; aryl; -NR₃₂R₃₃ in which either R₃₂ and R₃₃ represent, independently, the hydrogen atom, a lower alkyl, arylalkyl or alkylcarbonyl radical, or R₃₂ and R₃₃ form, with the nitrogen atom to which they are attached, a
25 heterocycloalkyl; or -Z₃₂-R₃₄ in which Z₃₂ represents O, C(O), OC(O), S, S(O) or SO₂ and R₃₄ represents the hydrogen atom, a lower alkyl, aryl or lower arylalkyl radical;

a -Z₃₁R₃₁ radical in which Z₃₁ represents O, C(O), OC(O), S, and R₃₁ represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical.


30 In the definitions indicated above, the expression halo represents the fluoro, chloro, bromo or iodo radical, preferably chloro, fluoro or bromo. The expression lower alkyl preferably represents a linear or branched alkyl radical having 1 to 6 carbon atoms, and in particular an alkyl radical having 1 to 4 carbon atoms such as the methyl, ethyl, propyl, isopropyl, butyl,

isobutyl, sec-butyl and tert-butyl radicals, but can also represent a pentyl, isopentyl, hexyl or isohexyl radical. Among the alkyl radicals containing 1 to 10 carbon atoms, the lower alkyls as defined above can be mentioned but also the heptyl, octyl, nonyl or decyl radicals.

5 The lower alkoxy radicals can correspond to the alkyl radicals indicated above such as for example the methoxy, ethoxy, propyloxy or isopropyloxy radicals but also linear, secondary or tertiary butoxy. The term lower alkylthio preferably designates the radicals in which the alkyl radical is as defined above such as for example methylthio, ethylthio.

10 The expression lower alkenyl preferably represents a linear or branched alkenyl radical having 1 to 6 carbon atoms, such as for example vinyl, allyl, propenyl, butenyl or pentenyl. The terme lower alkynyl preferably represents a linear or branched alkynyl containing 1 to 6 carbon atoms, and preferably an ethynyl, propargyl, butynyl or pentynyl radical.

15 The terme cycloalkyl preferably designates the cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl rings. The expression heterocycloalkyl designates a saturated cycloalkyl containing 2 to 7 carbon atoms and at least one heteroatom. This radical can contain several identical or different heteroatoms. Preferably, the heteroatoms are chosen from oxygen, sulphur or nitrogen. As examples of heterocycloalkyl, there can be mentioned the pyrrolidine, imidazolidine, pyrrazolidine, isothiazolidine, thiazolidine, isoxazolidine, oxazolidine ring or a ring of formula



20 in which E represents CH₂, O or R₄N in which R₄ represents hydrogen, a lower alkyl, arylalkyl, arylsulphonyl, optionally substituted aryl radical, thus being able to represent for example the piperidine, piperazine or morpholine ring.

25 The expression aryl represents an aromatic radical, constituted by a ring or condensed rings, such as for example the phenyl or naphthyl radical. The term aryloxy preferably designates the radicals in which the aryl radical is as defined above such as for example the phenoxy radical. The expression heteroaryl designates an aromatic radical, constituted by a ring or condensed rings, with at least one ring containing one or more identical or different heteroatoms chosen from sulphur, nitrogen or oxygen. As an example of a heteroaryl radical, thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, thiazolyl, isoxazolyl, oxazolyl, 30 pyridyl, pyrazinyl, pyrimidyl, benzothienyl, benzofuryl and indolyl radicals can be mentioned.

The lower arylalkyl radicals designate the radicals in which respectively the aryl and lower alkyl radicals are as defined above such as for example benzyl, phenethyl or naphthylmethyl. The lower heteroarylalkyl radicals designate the radicals in which respectively the heteroaryl

and lower alkyl radicals are as defined above such as for example indolylmethyl, thienylmethyl, furylmethyl. The term aryloxyalkyl designates the radicals in which respectively the aryloxy and lower alkyl radicals are as defined above.

5 The terms lower alkyl amino and lower dialkyl amino preferably designate the radicals in which the alkyl radicals are as defined above, such as for example methylamino, ethylamino, dimethylamino, diethylamino or (methyl)(ethyl)amino.

10 The terms alkylsulphonyl, cycloalkylsulphonyl, arylsulphonyl and arylalkylsulphonyl preferably designate the radicals in which respectively the alkyl, cycloalkyl and aryl radicals are as defined above. Similarly, the terms alkylcarbonyl, alkoxycarbonyl, arylcarbonyl, aryloxy carbonyl preferably designate the radicals in which respectively the alkyl, alkoxy, aryl and aryloxy radicals are as defined above.

When the products of formula I contain a basic group, they can form addition salts with acids, in particular pharmacologically acceptable acids.

15 According to the definition of the variable groups, a compound of formula I as defined above can have one or more asymmetrical carbons. The invention relates to the use of the compounds of formula I as defined above, which compounds can be found in racemic, enantiomeric or diastereoisomeric form.

20 A more particular subject of the invention is the use, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors are involved, of a compound of general formula I as defined above in which

W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

25 R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals: lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, aryl, aryloxy or heterocycloalkyl;

R_1 represents one or more identical or different groups, chosen from: the hydrogen atom, a halo, lower alkyl or lower alkoxy radical;

30 R_{2a} and R_{2b} represent, independently the hydrogen atom or a lower alkyl radical;

R₃ represents the hydrogen atom; an alkyl radical with 1 to 10 carbon atoms, cycloalkylalkyl, aryl, lower arylalkyl or lower heteroarylalkyl, the alkyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted by one or more identical or different radicals chosen from:

- 5 aryl; -NR₃₂R₃₃ in which either R₃₂ and R₃₃ represent, independently, the hydrogen atom or a lower alkyl radical; or -Z₃₂-R₃₄ in which Z₃₂ represents O and R₃₄ represents the hydrogen atom or a lower alkyl radical.

A more particular subject of the invention is the use, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved, of a compound of formula I as defined above,
10 characterized in that

W represents the hydrogen atom or a radical of formula R-X-C(Y)-;

R represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the
15 following radicals: methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino;

X represents CH₂, C₂H₄, CH₂NH, NH, O, S or a covalent bond;

20 Y represents O or S;

R₁ represents one or more identical or different groups, chosen from: the hydrogen atom, a chloro, methyl or methoxy radical;

R_{2a} and R_{2b} represent, independently, the hydrogen atom or a methyl radical;

R₃ represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy, phenethyl, naphthylmethyl or
25 indolylmethyl radical.

More particularly, a subject of the invention is the use, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved, of the compounds described hereafter in the examples, in
30 particular the products corresponding to the following formulae:

- 1-butyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 5 - 1-methyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-ethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 10 - 1-propyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-phenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 15 - 1-pentyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-hexyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-(4-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 20 - 1-(4-methoxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-(1-naphtyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 25 - 1-(3-indolyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-phenethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-diphenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-ethoxyethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 5 - 1-cyclohexylmethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-(3-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-(dimethylaminoethyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 10 - 1-methyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-benzyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 15 - 1-benzyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-methyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-benzyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 20 - 1-methyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-butyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 25 - 1-benzyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-butyl-6-(2-methoxyphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;


- 1-heptyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-hexyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 5 - 1-pentyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-oxoethyl]-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-thioxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 10 - 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide;
- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxamide;
- 15 - 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylbenzyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide;
- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-benzyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxamide;
- phenyl ester of 6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxylic acid;
- 20 - 6-(2-chlorophenyl)-7,10-dihydro-1,4-dimethyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide;
- 1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide;
- 25 but also of the products of formula I as defined above and of which the substituents R,X,Y, R₁, R_{2a}, R_{2b} and R₃ are respectively the following:
- 2-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-isoPr-Ph; NH; S; 2-Cl; H; H; Me;

- 2-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Et;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; H;
- 2-terBu-Ph; NH; S; 2-Cl; H; H; Me;
- 5 - 1-naphthyl; NH; S; 2-Cl; H; H; Me;
- 2-F₃CO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Et-Ph; NH; S; 2-Cl; H; H; Me;
- 10 - 2-PhO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Pr-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-EtOC(O)-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeS-Ph; NH; S; 2-Cl; H; H; Me;
- 15 - 2-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,4-(MeO)-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 20 - 2,3-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-5-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 25 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-5-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 30 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; Pr;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Bu;
- 2-F₃C-Ph; NH; S; H; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Pr;
- 35 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Bu;
- 2-MeSO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Bz;

- 2-F₃C-Ph; NH; S; 4-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; pentyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; hexyl;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Bz;
5 - 2-NO₂-4-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 1-naphthyl-methyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-indolyl-methyl;
- 2-MeS-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
10 - 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-EtO-Ph; NH; S; 2-Cl; H; H; Me;
15 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-MeO-Bz;
- 2-NO₂-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-HO-Bz;
- 2-F₃C-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
20 - 2-NO₂-4-MeO-Ph; NH; S; H; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-C₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; EtOC₂H₄;
- 3-NO₂-2-pyridinyl; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; -, O; 2-Cl; H; H; Me;
25 - Ph; -, S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Me;
- 2-NO₂-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bu;
30 - 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bu;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; cyclohexylmethyl;
35 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; (Me)₂NC₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-HO-Bz;
- Ph; S; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; heptyl,

as well as the salts of these compounds with mineral or organic acids when the molecule contains a basic group.

Among the compounds of formula I, some are new. A subject of the invention is therefore also a compound of general formula II



in which

W' represents the hydrogen atom or a radical of formula $R'-X'-C(Y')-$;

R' represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the following radicals: methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethyloxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino;

X' represents CH_2 , C_2H_4 , CH_2NH , NH , O , S or a covalent bond;

Y' represents O or S ;

R'_1 represents one or more identical or different groups, chosen from: the hydrogen atom, a chloro, methyl or methoxy radical;

$R_{2a'}$ and $R_{2b'}$ represent, independently, the hydrogen atom, a methyl radical;

R'_3 represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy radical, phenethyl, naphthylmethyl or indolylmethyl.

with the exclusion of the compounds of formula II in which

W' represents the hydrogen atom, R'₁ represents the chloro radical in ortho position; R_{2a'} represents the hydrogen atom and R_{2b'} represents the hydrogen atom or the methyl radical; R'₃ represents the methyl radical and

W' represents a radical of formula R'-X'-C(Y')- and

5 - X' represents NH; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents the 4-terbutylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 2,3-dichlorophenyl, 2,4-(difluoro)phenyl, 4-phenoxyphenyl; pyridyl; cyanophenyl radical;

10 - X' represents NH; Y' represents S; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents the 4-terbutylphenyl, 2,4-diterbutylphenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 4-fluorophenyl, 4-(methylsulphonyl)phenyl radical;

15 - X' represents CH₂NH; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl.

20 - X' represents the oxygen atom or a covalent bond; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents pyridyl or cyanophenyl;

 - X' represents CH₂; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl or 4-fluorophenyl;

25 - X' represents C₂H₄; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl.

More particularly, a subject of the invention is a compound of general formula II in which W' represents a radical of formula R'-X'-C(Y')- and the substituents R' , X' , Y' , R'₁ , R_{2a'} , R_{2b'} and R'₃ represent respectively:

30 - 2-F₃C-Ph; CH₂; O; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; NH; O; 2-Cl; H; H; Me;


- 2-F₃C-Ph; CH₂NH; S; 2-Cl; H; H; Me;
- Ph; O; O; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; Me; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Bz;
5 - 3-F₃C-Ph; NH; O; 2-Cl; H; H; Me;
- 4-F₃C-Ph; NH; O; 2-Cl; H; H; Me;
- 2-isoPr-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Et;
10 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; H;
- 2-terBu-Ph; NH; S; 2-Cl; H; H; Me;
- 1-naphthyl; NH; S; 2-Cl; H; H; Me;
- 2-Ph-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃CO-Ph; NH; S; 2-Cl; H; H; Me;
15 - 2-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Et-Ph; NH; S; 2-Cl; H; H; Me;
- 2-PhO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Pr-Ph; NH; S; 2-Cl; H; H; Me;
20 - 2-EtO-Ph; NH; S; 2-Cl; H; H; Me;
- Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-EtOC(O)-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeS-Ph; NH; S; 2-Cl; H; H; Me;
25 - 2-morpholino-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2,6-isoPr-Ph; NH; S; 2-Cl; H; H; Me;
- 2,6-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-(MeO)-Ph; NH; O; 2-Cl; H; H; Me;
30 - 2-MeO-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,4-(MeO)-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,3-Cl-Ph; NH; S; 2-Cl; H; H; Me;
35 - 2,5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-3-Cl-Ph; NH; S; 2-Cl; H; H; Me;

- 2-Me-5-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2,3-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
5 - 2-NO₂-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-5-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
10 - 2-Cl-5-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Pr;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Bu;
- 3-Ph-6-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; H; H; H; Me;
15 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Pr;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Bu;
- 2-NO₂-4-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeSO₂-Ph; NH; S; 2-Cl; H; H; Me;
20 - 2-F₃C-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Bz;
- 2-F₃C-Ph; NH; S; 4-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; pentyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; hexyl;
25 - 3,5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Bz;
- 2-NO₂-4-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 1-naphthyl-methyl;
30 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-indolyl-methyl;
- 2-MeS-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-HO-Ph; NH; S; 2-Cl; H; H; Me;
35 - 2-NO₂-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-EtO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-MeO-Bz;

- 2-NO₂-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-HO-Bz;
- 2-F₃C-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 5 - 2-NO₂-4-MeO-Ph; NH; S; H; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-C₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; EtOC₂H₄;
- 3-NO₂-2-pyridinyl; NH; S; 2-Cl; H; H; Me;
- 4-MeO-Ph; CH₂; O; 2-Cl; H; H; Me;
- 10 - 2-indolyl; -, O; 2-Cl; H; H; Me;
- 3-indolyl; CH₂; O; 2-Cl; H; H; Me;
- 4-HO-Ph; C₂H₄; O; 2-Cl; H; H; Me;
- 2-F₃C-Ph; -, O; 2-Cl; H; H; Me;
- 4-HO-Ph; CH₂; O; 2-Cl; H; H; Me;
- 15 - 5-MeO-2-indolyl; -, O; 2-Cl; H; H; Me;
- Ph; -, O; 2-Cl; H; H; Me;
- Ph; -, S; 2-Cl; H; H; Me;
- 5-MeO-2-indolyl; -, S; 2-Cl; H; H; Me;
- 2-NO₂-Ph; CH₂; O; 2-Cl; H; H; Me;
- 20 - 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Me;
- 2-NO₂-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bu;
- 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bz;
- 25 - 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bu;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; cyclohexyl methyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; (Me)₂NC₂H₄;
- 30 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-HO-Bz;
- 2-pyridinyl; NH; S; 2-Cl; H; H; Me;
- Ph; S; S; 2-Cl; H; H; Me;
- Ph; O; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; heptyl;
- 35 but also the compounds of formula II in which W' represents the hydrogen atom and the substituents R'₁, R'_{2a}, R'_{2b} and R'₃ represent respectively:
- 2-Cl; H; H; butyl;
- 2-Cl; H; H; benzyl;


- 2-Cl; H; H; H;
- 2-Cl; H; H; ethyl;
- 2-Cl; H; H; propyl;
- 2-Cl; H; H; Ph;
- 5 - 2-Cl; H; H; pentyl;
- 2-Cl; H; H; hexyl;
- 2-Cl; H; H; 4-HO-Bz;
- 2-Cl; H; H; 4-MeO-Bz;
- 2-Cl; H; H; 1-naphthyl-methyl;
- 10 - 2-Cl; H; H; 3-indolyl-methyl;
- 2-Cl; H; H; Ph-C₂H₄;
- 2-Cl; H; H; Ph-Ph;
- 2-Cl; H; H; EtOC₂H₄;
- 2-Cl; H; H; cyclohexylmethyl;
- 15 - 2-Cl; H; H; 3-OH-Bz;
- 2-Cl; H; H; (Me)₂NC₂H₄;
- H; H; H; Me;
- 4-Cl; H; H; Bz;
- H; H; H; Bz;
- 20 - 4-Cl; H; H; Me;
- 3-Cl; H; H; benzyl;
- 3-Cl; H; H; Me;
- 2-Me; H; H; butyl;
- 2-Me; H; H; benzyl;
- 25 - 2-MeO; H; H; butyl;
- 2-Cl; H; H; heptyl;
- 2-Cl; H; H; hexyl;
- 2-Cl; H; H; butyl.

30 The compounds of formula I according to the invention and in which W represents the R-X-C(Y)- radical, can be prepared according to the process which consists of reacting a compound of formula (1)




in which R_1 , R_{2a} , R_{2b} and R_3 have the meaning indicated above, with, according to the final product chosen

A) either a compound of formula (2)




5 in which R , Y and n have the meaning indicated above, in order to form a compound of formula I in which X represents the $-(CH_2)_n NH$ - radical;

B) or a compound of formula (3)



10 in which R , Y and n have the meaning indicated above, Y'' represents O or S and A represents a halogen atom, in order to form a compound of formula I in which X represents the $-(CH_2)_n-O$ - or $-(CH_2)_n-S$ - radical;

C) or a compound of formula (4)




15 in which R and n have the meaning indicated above, in order to form a compound of formula I in which X represents the $-(CH_2)_n$ radical and Y the oxygen atom, which compound can be optionally converted to a compound of formula I in which X represents the $-(CH_2)_n$ radical and Y the sulphur atom.

During the preparation according to route A), the addition of the compound of formula (2) as defined above, with a compound of formula (1) is easily carried out at a temperature close to 20°C in a chlorinated solvent such as dichloromethane or 1,2-dichloroethane.


During the preparation according to route B), the conversion of compound (1) to a compound of formula I in which X represents the $-(CH_2)_nY^{\bullet}$ radical, by reaction with a compound of formula (3), can be carried out at a temperature close to 20°C in an inert solvent such as dichloromethane, and preferably in the presence of an acid acceptor such as triethylamine.

During the preparation according to route C), the compound of formula (1) is converted to a compound of formula I in which X represents the $-(CH_2)_n$ radical and Y the oxygen atom, by reaction with acid (4) under activation conditions similar to peptide coupling reactions. The reaction can be carried out at a temperature close to 20°C, in an inert solvent such as dimethylformamide, tetrahydrofuran or dichloromethane, and generally in the presence of an acid acceptor such as a tertiary amine such as for example, triethylamine or diisopropylethylamine. The amide thus obtained, can be converted to a compound of formula I in which X represents the $-(CH_2)_n$ radical and Y the sulphur atom, while heating the reaction medium at a temperature close to 90°C in a polar solvent such as toluene or cyclohexane, in the presence of a thiation agent such as Lawesson reagent.


The starting product of formula (1) corresponds to the product of formula I in which W represents a hydrogen atom. Such a compound can be obtained by reacting a compound of formula (5)



in which R_1 , R_{2a} and R_{2b} have the meaning indicated above and R'' represents a lower alkyl or lower arylalkyl radical, with a compound of formula (6)




in which R₃ has the meaning indicated above, in order to obtain compound (7)




which compound is then subjected to a deprotection reaction of the carbamate in order to obtain product (1).

- 5 The compound of formula (1) can also be prepared by reacting a compound of formula (5) as defined above with hydrazine in order to obtain a compound of formula (8)



which compound is then reacted with a trialkyl orthoalkanoate type compound of formula (9)



- 10 in which R₈ represents an alkyl group and R₃ has the meaning indicated above, in order to produce the compound of formula (7) as defined above which is then deprotected.

According to the first method for the preparation of compound 1, the compounds of formula (5) react on the compounds of formula (6) such as phenyl acethydrazide, 4-hydroxyphenyl acethydrazide, while heating an inert solvent such as toluene under reflux and in certain cases
 15 for reasons of solubility a mixture of polar solvents under reflux such as the

1,2-dichloroethane/methanol mixture. Preferably, compound (5) used is such that R" represents the ethyl or ter-butyl radical. The reaction is carried out in two stages with, in a first stage, formation of the corresponding azide followed by an intramolecular cyclization reaction leading to the compound of formula (7).

- 5 Deprotection of the carbamate can be carried out while heating in the presence of a mineral base such as sodium hydroxide in a lower aliphatic alcohol such as ethanol or by agitating at ambient temperature in highly acid medium such as for example hydrobromic acid (33% in acetic acid) in order to produce the compounds of formula (1). Other methods of cleaving the carbamate such as those described in Protective Groups in Organic Synthesis [T.W. Green,
10 P.G.M. Wuts; 2nd Edition, J. Wiley and sons Inc., p. 364-6 (1991)] can also be used in so far as they are compatible with the substituents constituting the pyrido thieno triazolo diazepine skeleton of the compounds of general formula (7).

- According to the second method, the compound of formula (5) reacts with hydrazine in solution in water while heating an aliphatic alcohol such as ethanol, isopropanol under reflux.
15 This intermediate hydrazine compound is then reacted with a trialkyl orthoalkanoate compound of formula (9) in order to produce a compound of formula (7).

The product of formula (5) can be prepared according to the method described in patent FR2645153 or according to similar methods.

- Certain products of formula (6) are commercially available; the others can be prepared by
20 reacting under reflux of ethanol, the alkyl ester of $R_3\text{-CO}_2\text{H}$ acid on the hydrazine diluted to 35% in water. The reagents (9), (4) and (3) are in general marketed (for example by the firm Acros or Aldrich). The products of formula $R\text{-(CH}_2)_n\text{-N=C=Y}$ are mostly commercially available or can be prepared by reacting the corresponding amine on (thio)phosgene according to methods known to a person skilled in the art.

- 25 In the case where R_{2a} and R_{2b} do not represent the hydrogen atom, the product of formula (1) can also be prepared in a similar fashion to that described in the Patent FR 2645153 with, for example, the use of 2-bromopropionyl bromide instead of bromoacetyl bromide in the second stage of the synthesis of the tetracycle.

- Compounds I of the present invention have useful pharmacological properties. Thus it has
30 been discovered that compounds I of the present invention have a high affinity for one (or more) of the somatostatin receptors. They can be used as non-peptide agonists or antagonists of somatostatin in a selective or non-selective manner.

The compounds of the present invention can therefore be used in different therapeutic applications. The compounds can advantageously be used to treat the pathological states or the diseases as presented above and in which one (or more) of the somatostatin receptors are involved.

- 5 An illustration of the pharmacological properties of the compounds of the invention will be found hereafter in the experimental part.

These properties make the products of formula I suitable for a pharmaceutical use. A subject of the present application is also, as medicaments, the products of formula II as defined above, as well as the addition salts with pharmaceutically acceptable mineral or organic acids of said
10 products of formula II, as well as the pharmaceutical compositions containing, as active ingredient, at least one of the medicaments as defined above.

The invention therefore relates to the pharmaceutical compositions containing a compound of formula II according to the invention or a pharmaceutically acceptable addition salt of acid of this, in combination with a pharmaceutically acceptable support. The pharmaceutical
15 composition can be in the form of a solid, for example, powders, granules, tablets, gelatin capsules or suppositories. Appropriate solid supports can be, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone and wax.

The pharmaceutical compositions containing a compound of the invention can also be presented
20 in liquid form, for example, solutions, emulsions, suspensions or syrups. Appropriate liquid supports can be, for example, water, organic solvents such as glycerol or glycols, similarly their mixtures, in varying proportions, in water, with added pharmaceutically acceptable oils or fats. The sterile liquid compositions can be used for intramuscular, intraperitoneal or subcutaneous injections and the sterile compositions can also be administered intravenously.

- 25 The following examples are presented to illustrate the above procedures and should in no event be considered as a limit to the scope of the invention.

EXPERIMENTAL PART:

Example 1

1-butyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4]
30 triazolo [4,3-a] [1,4] diazepine

1st stage

ethyl 5-(2-chlorophenyl)-2-hydrazino-6,9-dihydro-3H-pyrido[4',3';4,5]thieno[2,3-e][1,4]-diazepine- 8 (7H) carboxylate

Hydrazine at 35% in water (42 ml) is added at ambient temperature and under argon to a mixture containing 5-(2-chlorophenyl)-8-(ethoxycarbonyl)-6,7,8,9-tetrahydro-3H-pyrido
5 [4',3':4,5] thieno [3,2-e] [1,4] diazepine 2-thione (118.5 g, 0.282 mol) in 1740 ml of methanol. Agitation is maintained at this temperature for one hour. The precipitate is filtered on frit, washed with ethanol (100ml) and with ether (2x200ml) then dried under vacuum (61.9g). The solvent contained in the filtrate is evaporated off. The solid is filtered then washed with methanol and with ether (44.1g). The desired product is in the form of a yellow
10 solid (total yield: 106g, 90%). Melting point: 216°C.
NMR¹H (100 MHz, DMSO d₆, δ): 1.15 (t, 3H); 1.62 (m, 2H); 3.34 (m, 2H); 4.00 (q, 2H); 4.40-4.60 (m, 4H); 7.34-7.41 (m, 4H); 9.13 (m, 1H)
IR (cm⁻¹): ν_{NH} (hydrazine): 2850-2950; ν_{C=O} (carbamate): 1690; 1600; 1310; 1370; 1240; 1120; 1050; 760.

15 2nd stage

ethyl 1-butyl-6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a][1,4] diazepine-9(8H)-carboxylate

A mixture containing ethyl 5-(2-chlorophenyl)-2-hydrazino-6,9-dihydro-3H-pyrido[4',3';4,5] thieno [2,3-e] [1,4]-diazepine-8(7H) carboxylate (3g, 7.2 mmol) and trimethyl orthovalerate
20 (1.2 ml, 7.2 mol) in 15 ml of butanol is heated under reflux. Agitation is carried out for twelve hours at 20°C until precipitation of the product. The latter is filtered on frit, washed with ethanol (15 ml), with isopropanol (15 ml), with isopropyl ether (15 ml) and with isopentane (15 ml) then dried under vacuum in order to obtain a white solid (2g, 58%).
Melting point: 190°C
25 NMR¹H (400 MHz, DMSO d₆, δ): 0.85 (t, 3H); 1.14 (m, 3H); 1.3 (m, 2H); 1.47 (m, 1H); 1.61 (m, 2H); 2.10 (d, 1H); 3.00 (t, 2H); 3.11 (m, 1H); 3.71 (m, 1H); 4.02 (q, 2H); 4.24 (d, 1H); 4.43 (m, 1H); 4.82 (d, 1H); 5.32 (d, 1H); 7.43-7.76 (m, 4H)
IR (cm⁻¹): ν_{C=O} (carbamate): 1690; 1606; 1438; 1417; 1230; 1121; 761.

3rd stage

30 1-butyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

A mixture containing ethyl 1-butyl-6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno[3,2-f] [1,2,4] triazolo [4,3a] [1,4] diazepine-9(8H)-carboxylate (1.98g, 4.8 mmol) and
35 potassium hydroxide (3g, 0.052 mol) in 40 ml of ethanol is heated under reflux of ethanol and

for ten hours. The reaction medium is cooled down to 22°C then filtered on frit. The frit is rinsed with 5 ml of ethanol. 80 ml of water is added to the filtrate. Five drops of acetic acid are added and the aqueous phase is extracted with dichloromethane (2x80 ml). The organic phase is dried over magnesium sulphate. The solvent is evaporated off with a rotary
5 evaporator. The reaction mixture is agitated for one hour in a solvent mixture of isopropyl ether-isopentane-isopropanol (10-10-1 ml), followed by filtering on frit then washing with isopropyl ether and with isopentane. After drying under vacuum a white solid is obtained (0.9g, 46%). Melting point: 224°C.

NMR¹H (400 MHz, DMSO d₆, δ): 0.85 (t, 3H); 1.28-1.40 (m, 3H); 1.59 (m, 2H); 1.94 (m,
10 1H); 2.55 (m, 1H); 2.78 (m, 1H); 2.99 (t, 2H); 3.84 (q, 2H); 4.22 (d, 1H); 5.30 (d, 1H); 7.43-7.47 (m, 4H)

IR (cm⁻¹): ν_{NH}: 3268; 2930; 1600; 1430; 849; 747.

Example 2

1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4]
15 triazolo [4,3-a] [1,4] diazepine

1st stage

Ethyl ester of 1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f]
[1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxylic acid

A mixture containing 5-(2-chlorobenzoyl)-8-(ethoxycarbonyl)-6,7,8,9-tetrahydro-3H-pyrido
20 [4',3';4,5] thieno [3,2 - f] [1,4] diazepine 2-thione (3 g, 0.007 mol) and the hydrazide of phenylacetic acid (1.18 g, 0.0077 mol) is heated under reflux in anhydrous toluene (40 ml) for 3 hours. After cooling down to ambient temperature, the precipitate is filtered on frit then agitated in a mixture of solvent (isopropyl ether/dichloromethane/isopentane: 90/8/2) for 12 hours. The solid is filtered on frit then washed with isopropyl ether, with isopropanol then
25 with isopentane. After drying under vacuum for 12 hours, 2.8 g (76 %) of a beige coloured solid is obtained. Melting point: 192°C.

NMR¹H (400MHz CDCl₃, δ): 1.26 (t, 3H); 1.63-2.03 (m, 2H); 3.12 (m, 1H); 3.85 (m, 1H);
4.18 (m, 3H); 4.45 (s, 1H); 4.47 (q, 2H); 4.82 (m, 1H); 5.61 (d, 1H); 7.17-7.34 (m, 9H).

IR (cm⁻¹): 1694 νC=O(carbamate); 1434; 1420; 1232; 1126; 758; 727.

2nd stage

1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4]
triazolo [4,3-a] [1,4] diazepine

A mixture containing ethyl ester of 1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-4H-
35 pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxylic acid

(3.69 g, 0.0071 mol) and potassium hydroxide (4.4 g, 0.078 mol) is heated under reflux in ethanol for 6 hours then left under agitation at ambient temperature for 48 hours. The solid is filtered on frit, washed with a minimum quantity of ethanol. After adding water, extraction is carried out with dichloromethane followed by drying over magnesium sulphate. The solvent
5 is evaporated off and the product is purified by crystallization from an ethyl acetate/isopropanol 50/50 mixture, followed by leaving in a refrigerator for 48 hours then filtering on frit while washing with a solvent mixture: ethyl acetate-isopropanol (50-50). Washing is finally carried out with isopentane followed by drying under vacuum. 2.46 g (77%) of the desired product is obtained in the form of a beige coloured solid. Melting point:
10 191-192°C.

NMR¹H (400MHz, CDCl₃, δ): 1.30 (m, 1H); 1.89 (m, 1H); 2.51 (m, 1H); 2.50 (se, 1H); 2.76 (m, 1H); 3.82 (q, 2H); 4.23 (dd, 1H); 4.44 (m, 2H); 5.33 (dd, 1H); 7.13-7.44 (m, 9H)
IR (cm⁻¹): 3422, 2928, 1602, 1427, 1419, 1045, 1033, 759, 725, 695.

Example 3

15 1-methyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point: 202°C.

Example 4

20 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point: 162°C.

Example 5

1-ethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

25 It is obtained according to the method as described in Example 1. Melting point: 242°C.

Example 6

1-propyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point > 260°C.

30 **Example 7**

1-phenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 262°C

Example 8

5 1-pentyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 174°C

Example 9

10 1-hexyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 194-198°C

Example 10

1-(4-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

15 It is obtained according to the method as described in Example 2. Melting point = 220-230°C

Example 11

1-(4-methoxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 254-256°C

20 **Example 12**

1-(1-naphthyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 185-190°C

Example 13

25 1-(3-indolyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 196-200°C

Example 14

1-phenethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 216-218°C

Example 15

5 1-diphenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 210-214°C

Example 16

10 1-ethoxyethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 152°C

Example 17

1-cyclohexylmethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

15 It is obtained according to the method as described in Example 2. Melting point = 188-190°C

Example 18

1-(3-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 242-246°C

20 **Example 19**

1-(dimethylaminoethyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 143-147°C

Example 20

25 1-methyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 240-244°C

Example 21

1-benzyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 238-242°C

Example 22

- 5 1-benzyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 214-218°C

Example 23

- 10 1-methyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 262°C

Example 24

1-benzyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

- 15 It is obtained according to the method as described in Example 2. Melting point = 170°C

Example 25

1-methyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 160°C

- 20 **Example 26**

1-butyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 206°C

Example 27

- 25 1-benzyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 2. Melting point = 110-120°C

Example 28

1-butyl-6-(2-methoxyphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 190°C

Example 29

5 1-heptyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 178-180°C

Example 30

10 1-hexyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

It is obtained according to the method as described in Example 1. Melting point = 190-192°C

Example 31

1-pentyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

15 It is obtained according to the method as described in Example 1. Melting point = 250°C

Example 32

6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-oxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

20 A mixture containing 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] 1,2,4-triazolo [4,3-a] 1,4-diazepine (2.22g, 0.006 mol), 2-trifluoromethylphenyl acetic acid (1.35g, 0.0066 mol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.26g, 0.0066 mol) hydrochloride and triethylamine (1 ml, 0.0066 mol) in 40 ml of anhydrous dimethylformamide is agitated at a temperature close to 20°C for 72 hours. The solvent is evaporated off with a rotary evaporator while entraining it with toluene. The crude reaction product is taken up in 30 ml of dichloromethane, washed with water and the organic phase is dried over magnesium sulphate and concentrated with a rotary evaporator. Then approximately 4 ml of dichloromethane is added to the reaction mixture then ether is added until precipitation. Agitation is carried out for a few minutes then filtering under vacuum while washing the precipitate with ether. The product obtained is dried under vacuum in order to obtain 2g (61%) of expected compound in the form of a white solid. Melting point: 224-226°C

25
30

HPLC- Conditions:

column: kromasyl C18 (150 x 4.6mm)

Mobile phase: Solution A: 0.02% trifluoroacetic acid

Solution B: acetonitrile

Gradient:	Temps (min.)	%A	%B	Curve
	0	90	10	linear
	5	90	10	
	14	30	70	
	15	10	90	
	25	10	90	
	25.5	90	10	
	55	90	10	

5

Retention time (min.): 14.82

Mass: MH^+ (exp.): 556.1; MH^+ (theoretical): 556.12

Example 33

6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-

10 1-thioxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine

A mixture containing 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-oxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (1g, 0.0018 mol) and Lawesson reagent (0.73g, 0.0018 mol) is heated for 2 hours at a temperature close to 80°C. Agitation is maintained for 14 hours at a temperature close to 20°C. The precipitate is filtered on frit, washed with 30 ml of toluene and 30 ml of acetone, dried under vacuum. The expected product is isolated by flash chromatography on a silica column with a dichloromethane-methanol mixture (95-5) as eluent

15

White solid. Melting point: >260°C

HPLC- Conditions:

20

column: kromasyl C18 (150 x 4.6mm)

Mobile phase: Solution A: 0.02% trifluoro acetic acid

Solution B: acetonitrile

Gradient:	Temps (min.)	%A	%B	Curve
	0	90	10	linear
	5	90	10	
	25	10	90	
	30	10	90	
	31	90	10	
	60	90	10	

Retention time (min.): 17.3

Mass: MH^+ (exp.): 572.1; MH^+ (theoretical): 572.09

Example 34

5 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide

1st stage

2-trifluoromethylphenylisothiocyanate

2-trifluoromethylaniline (4.1 ml, 0.0325 mol) is added dropwise at a temperature close to 20°C and under an argon atmosphere to a mixture containing thiophosgene (2.5 ml, 0.0326 mol) in 10 43 ml of water. Agitation is carried out for twelve hours then 30 ml of ethyl acetate is added. The organic phase is extracted with water then with a solution of 10% sodium bicarbonate and finally with a solution of salt water. The organic phase is dried over magnesium sulphate and the solvent is evaporated off with a rotary evaporator. The product obtained (6.85g, 100%) is in the form of a brown oil which is used immediately for the following reaction.

15 2nd stage

6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide

A mixture containing 2-trifluoromethylphenyl isothiocyanate (0.82g, 0.004mol) and 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (1g, 0.0027 mol) in 30 ml of 1,2-dichloroethane is agitated for 4 20 hours and at a temperature close to 20°C. Half the solvent is evaporated off with the rotary evaporator then 10 ml of ether is added. Agitation is carried out for 20 minutes at a temperature close to 20°C followed by filtering on frit then washing with 20 ml of ether, with 20 ml of isopropyl ether and with 20 ml of isopentane. After drying under vacuum a cream coloured 25 solid is obtained (1.21g, 78%). Melting point: 190-192°C

NMR 1H (400 MHz, DMSO d_6 , δ): 1.72 (m, 1H); 2.23 (m, 1H); 2.62 (s, 3H); 3.40 (m, 1H); 4.30 (m, 2H); 4.84 (d, 1H); 5.35 (d, 1H); 5.68 (d, 1H); 7.31-7.70 (m, 8H); 9.30 (s, 1H).

IR: $\nu_{NH}(\text{thiourea})$: 3250 cm^{-1} ; 1607 cm^{-1} ; 1525 cm^{-1} ; 1381 cm^{-1} ; 1317 cm^{-1} ; 757 cm^{-1}

Example 35

30 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxamide

A mixture containing 2-trifluoromethylphenyl isocyanate (1.5g, 0.008 mol) and 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (2.1g, 0.0057 mol) in 50 ml of 1,2-dichloroethane is agitated at ambient temperature for 2 hours. The reaction mixture is evaporated to dryness. The residual oil is precipitated by adding diisopropyl acetate. After filtration on frit, washing is carried out with ether and the solid is dried at 65°C overnight (2.2g, 69%). A beige powder is obtained. Melting point: 196-198°C.

NMR¹H (400 MHz, DMSO d₆, δ): 1.63 (m, 1H); 2.12 (m, 1H); 2.61 (s, 3H); 3.20 (m, 1H); 3.90 (m, 1H); 4.27 (d, 1H); 4.38 (d, 1H); 4.98 (d, 1H); 5.34 (d, 1H); 7.32-7.68 (m, 8H); 8.39 (s, 1H)
IR: ν_{NH}(urea): 3333 cm⁻¹; ν_{C=O} (carbonyl urea): 1670 cm⁻¹; 1520 cm⁻¹; 1318 cm⁻¹; 1112 cm⁻¹; 764 cm⁻¹.

Example 36

6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylbenzyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9 (8H)-carbothioamide

1st stage

2-trifluoromethylbenzylisothiocyanate

A solution containing 2-trifluoromethylbenzylamine (3.50g, 0.02 mol) in 50 ml of dichloromethane and a solution containing sodium bicarbonate (3.4g, 0.04 mol) in 55 ml of water are added simultaneously over one hour at a temperature close to 20°C to a solution of thiophosgene (2.28g, 0.02 mol) in 15 ml of dichloromethane. Agitation is carried out for 30 minutes. The reaction mixture is decanted and the organic phase washed with 100 ml of water then with 100 ml of a solution of salt water. The organic phase is dried over magnesium sulphate and the solvent is evaporated off with a rotary evaporator. The expected compound is obtained in the form of an oil (3.50g, 81%). Considering its reactivity, it is used immediately for the following reaction.

2nd stage

6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylbenzyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9 (8H)-carbothioamide

A mixture containing 2-trifluoromethylbenzylisothiocyanate (1g, 0.0046 mol) and 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (1.22g, 0.0033 mol) in 20 ml of dichloromethane is agitated at a temperature close to 20°C for 1 hour. The reaction mixture is evaporated to dryness. 40 ml of ethyl acetate is added to the residual oil. After separation of the insoluble part, the solvent is evaporated off. 40 ml of isopropyl ether is added until precipitation of the product then

filtration on frit and washing with isopropyl ether. The solid obtained (1.2g, 63%) is dried and a beige powder is obtained. Melting point: 178-184°C.

NMR¹H (400 MHz, DMSO d₆, δ): 1.68 (m, 1H); 2.21 (m, 1H); 2.67 (s, 3H); 3.49 (m, 1H); 4.20 (m, 2H); 4.86 (m, 1H); 5.11 (m, 2H); 5.38 (m, 1H); 5.60 (m, 1H); 6.03 (t, 1H); 7.35-7.71 (m, 8H).

IR: νNH(thiourea): 3266 cm⁻¹; 1607 cm⁻¹; 1313 cm⁻¹; 1117 cm⁻¹; 761 cm⁻¹

Example 37

6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-benzyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxamide

10 A mixture containing benzylisocyanate (133g, 0.01 mol) and 6-(2-chlorophenyl)- 7,8,9,10 tetrahydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (2.956g, 0.008 mol) in 50 ml of benzene is agitated at ambient temperature for 2 hours and 30 minutes. The precipitate is filtered then washed with ether then crystallized from a solvent mixture: ethanol-isopropyl alcohol, followed by filtering warm then the reaction medium is
15 left to crystallize cold overnight. After filtration of the crystals on frit, washing is carried out with isopropanol then with ether, followed by drying under vacuum for approximately forty hours at a temperature close to 75°C (2.5g, 62.5%). Melting point: 146-149°C.

NMR¹H (400 MHz, DMSO d₆, δ): 1.77 (m, 1H); 2.11 (m, 1H); 2.66 (s, 3H); 3.19 (m, 1H); 3.73 (m, 1H); 4.21 (d, 1H); 4.43 (t, 2H); 4.49 (d, 1H); 4.86 (d, 1H); 4.96 (t, 1H); 5.58 (d, 1H);
20 7.27-7.35 (m, 9H).

IR: νNH (urea): 3302 cm⁻¹; νC=O (carbonyl urea): 1653 cm⁻¹; 1550 cm⁻¹; 1268 cm⁻¹; 762 cm⁻¹; 724 cm⁻¹

Example 38

phenyl ester of 6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4]
25 triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxylic acid

Phenylchloroformate (0.19 ml, 0.0015 mol) and triethylamine (0.2 ml, 0.0015 mol) are added dropwise at 0°C and under an inert atmosphere to a mixture containing 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (0.37g, 0.001 mol) in 5 ml of dichloromethane. Agitation is carried out at ambient
30 temperature for two hours. The organic phase is washed with water and dried over magnesium sulphate. The solvent is evaporated with a rotary evaporator, followed by taking up in ether until precipitation. Purification is carried out by flash chromatography on a silica column with dichloromethane-methanol: 95-5 as eluent and the column filtrate is concentrated

with the rotary evaporator, followed by taking up in ether until a white precipitate (0.15g, 31%) is obtained.

Melting point: 218-220°C

HPLC- Conditions:

- 5 column: kromasyl C18 (150 x 4.6mm)
Mobile phase: Solution A: 0.02%trifluoro acetic acid
Solution B: acetonitrile

Gradient:	Temps (min.)	%A	%B	Curve
	5	90	10	linear
	25	30	70	
	30	30	70	
	31	90	10	
	45	90	10	

- 10 Retention time (min.): 14.82
Mass: MH⁺ (exp.): 490.07; MH⁺(theoretical): 490.11

Example 39

6-(2-chlorophenyl)-7,10-dihydro-1,4-dimethyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide

- 15 A mixture containing 2-trifluoromethylphenylisothiocyanate (0.3 ml, 0.0015 mol) and 6-(2-chlorophenyl)- 7,8,9,10 tetrahydro-1,4-dimethyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (0.383 mg, 0.001 mol) in 10 ml of anhydrous dichloromethane is agitated for two hours at a temperature close to 20°C. The reaction mixture is evaporated to dryness. Diisopropyl acetate is added until precipitation. After filtration of the insoluble part,
20 evaporation to dryness is carried out. Ether is added until precipitation. The precipitate is filtered on frit and washed with ether then dried under vacuum (0.35g, 59%) which produces a clear yellow powder. Melting point: 210-213°C

NMR¹H (400 MHz, DMSO d₆, δ): 1.73 (m, 1H); 1.89 (d,3H); 2.25 (d, 1H); 2.61 (s, 3H); 3.55 (m, 1H); 4.29-4.38 (m, 2H); 4.84 (d, 1H); 5.70 (d, 1H); 7.31-7.70 (m, 8H); 9.34 (s, 1H).

- 25 IR: νNH (thiourea): 3408 cm⁻¹; 1525 cm⁻¹; 1319 cm⁻¹; 759 cm⁻¹

Example 40

1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide

Trifluoromethylphenylisothiocyanate (0.56 g, 0.0027 mol) is added to a mixture containing 1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine (0.8g, 0.0018 mol) in anhydrous 1,2-dichloroethane (13 ml). The reaction medium is agitated at ambient temperature for 12 hours. The precipitate is filtered on frit then washed with isopropyl ether, with ethyl acetate then with isopentane and finally dried under vacuum. 0.28g (24 %) of final product is obtained in the form of a pale yellow coloured solid. Melting point: 239-240°C.

NMR¹H (400 MHz, CDCl₃, δ): 1.72 (m,1H); 2.15 (m,1H); 3.50 (m, 1H); 4.12 (m, 1H); 4.19 (d, 1H); 4.46 (q, 2H); 4.74 (d, 1H); 5.33 (d, 1H); 5.62 (d, 1H); 7.16-7.67 (m, 14H).

IR (cm⁻¹): 3349, 1527, 1318, 1128, 756.

HPLC (UV): 95.8 %

The products described in the table below which constitute Examples 41 to 148, are obtained according to the same processes as those described in Examples 32 to 40. The symbols used in the table are as follows: Me = methyl, Et = ethyl, Pr = propyl, Bu = butyl, Ph = phenyl, Bz = benzyl, Mp = melting point.

Ex	R	X	Y	R ₁	R _{2a}	R _{2b}	R ₃	Mp (°C)
41	2-MeO-Ph	NH	S	2-Cl	H	H	Me	165
42	2-Me-Ph	NH	S	2-Cl	H	H	Me	165
43	3-F ₃ C-Ph	NH	O	2-Cl	H	H	Me	222-223
44	4-F ₃ C-Ph	NH	O	2-Cl	H	H	Me	243-245
45	2-isoPr-Ph	NH	S	2-Cl	H	H	Me	212-213
46	2-NC-Ph	NH	S	2-Cl	H	H	Me	232-236
47	2-F ₃ C-Ph	NH	S	2-Cl	H	H	Et	225-230
48	2-F ₃ C-Ph	NH	S	2-Cl	H	H	H	250-255
49	2-terBu-Ph	NH	S	2-Cl	H	H	Me	180-182
50	1-naphthyl	NH	S	2-Cl	H	H	Me	194-196
51	2-Ph-Ph	NH	S	2-Cl	H	H	Me	165-170
52	2-F ₃ CO-Ph	NH	S	2-Cl	H	H	Me	220-223
53	2-Cl-Ph	NH	S	2-Cl	H	H	Me	200-203
54	2-F-Ph	NH	S	2-Cl	H	H	Me	206-208
55	2-Et-Ph	NH	S	2-Cl	H	H	Me	178-180
56	2-PhO-Ph	NH	S	2-Cl	H	H	Me	230-232
57	2-Pr-Ph	NH	S	2-Cl	H	H	Me	190-195
58	2-EtO-Ph	NH	S	2-Cl	H	H	Me	204-206
59	Ph	NH	S	2-Cl	H	H	Me	206-210

60	2-Br-Ph	NH	S	2-Cl	H	H	Me	190-195
61	2-EtOC(O)-Ph	NH	S	2-Cl	H	H	Me	166-170
62	2-MeS-Ph	NH	S	2-Cl	H	H	Me	178-180
63	2-morpholino-Ph	NH	S	2-Cl	H	H	Me	242-246
64	2-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	190-192
65	2,6-isoPr-Ph	NH	S	2-Cl	H	H	Me	160-165
66	2,6-Me-Ph	NH	S	2-Cl	H	H	Me	198-201
67	2,5-(MeO)-Ph	NH	O	2-Cl	H	H	Me	258-260
68	2-MeO-5-Cl-Ph	NH	S	2-Cl	H	H	Me	203-206
69	2,4-(MeO)-Ph	NH	S	2-Cl	H	H	Me	194-196
70	2-Cl-5-F ₃ C-Ph	NH	S	2-Cl	H	H	Me	218-220
71	2-Me-5-Cl-Ph	NH	S	2-Cl	H	H	Me	236-240
72	2,3-Cl-Ph	NH	S	2-Cl	H	H	Me	230-232
73	2,5-Me-Ph	NH	S	2-Cl	H	H	Me	192-196
74	2,5-Cl-Ph	NH	S	2-Cl	H	H	Me	234-236
75	2-Cl-4-Me-Ph	NH	S	2-Cl	H	H	Me	206-208
76	2-Me-3-Cl-Ph	NH	S	2-Cl	H	H	Me	248-252
77	2-Me-5-F-Ph	NH	S	2-Cl	H	H	Me	202-204
78	2,3-Me-Ph	NH	S	2-Cl	H	H	Me	225-228
79	2-F ₃ C-4Br-Ph	NH	S	2-Cl	H	H	Me	212-214
80	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Me	248-250
81	2-NO ₂ -4-Me-Ph	NH	S	2-Cl	H	H	Me	262-264
82	2-MeO-4-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	230-233
83	2,5-Br-Ph	NH	S	2-Cl	H	H	Me	260-264
84	2-MeO-5-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	205-208
85	2-Cl-4-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	244-248
86	2-Cl-5-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	252-254
87	2-F ₃ C-Ph	NH	S	2-Cl	H	H	Pr	229-230
88	2-F ₃ C-Ph	NH	S	2-Cl	H	H	Bu	216
89	3-Ph-6-MeO-Ph	NH	S	2-Cl	H	H	Me	260-262
90	2-F ₃ C-Ph	NH	S	-	H	H	Me	178-185
91	2-F ₃ C-Ph	NH	S	2-Cl	H	H	Ph	190-194
92	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Pr	255-260
93	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Bu	197-200
94	2-NO ₂ -4-F ₃ C-Ph	NH	S	2-Cl	H	H	Me	215-220
95	2-MeSO ₂ -Ph	NH	S	2-Cl	H	H	Me	190-195
96	2-F ₃ C-4-Cl-Ph	NH	S	2-Cl	H	H	Me	204-206

97	2-NO ₂ -4-MeO-Ph	NH	S	4-Cl	H	H	Bz	170-179
98	2-F ₃ C-Ph	NH	S	4-Cl	H	H	Me	192-196
99	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	pentyl	202-204
100	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	hexyl	103-105
101	3,5-F ₃ C-Ph	NH	S	2-Cl	H	H	Me	>260
102	2-NO ₂ -4-MeO-Ph	NH	S	3-Cl	H	H	Bz	165-170
103	2-NO ₂ -4-F-Ph	NH	S	2-Cl	H	H	Me	176-180
104	2-NO ₂ -4-NC-Ph	NH	S	2-Cl	H	H	Me	206-210
105	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	1-naphthyl -methyl	215-218
106	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	3-indolyl -methyl	231-235
107	2-MeS-5-F ₃ C-Ph	NH	S	2-Cl	H	H	Me	185-189
108	2-NO ₂ -4-MeO-Ph	NH	S	3-Cl	H	H	Me	192-196
109	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Me	212-214
110	2-NO ₂ -4-HO-Ph	NH	S	2-Cl	H	H	Me	225-230
111	2-NO ₂ -5-Cl-Ph	NH	S	2-Cl	H	H	Me	196-200
112	2-NO ₂ -5-Me-Ph	NH	S	2-Cl	H	H	Me	202-204
113	2-NO ₂ -4-EtO-Ph	NH	S	2-Cl	H	H	Me	213-215
114	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	4-MeO- Bz	178-180
115	2-NO ₂ -4-Cl-Ph	NH	S	2-Cl	H	H	Me	226-230
116	2-Br-4-Me-Ph	NH	S	2-Cl	H	H	Me	178-180
117	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	4-HO-Bz	196-197
118	2-F ₃ C-4-NO ₂ -Ph	NH	S	2-Cl	H	H	Me	175-180
119	2-NO ₂ -4-MeO-Ph	NH	S	H	H	H	Bz	201-202
120	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Ph-C ₂ H ₄	165-170
121	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	EtOC ₂ H ₄	145-153
122	3-NO ₂ -pyridyl	NH	S	2-Cl	H	H	Me	224-226
123	4-MeO-Ph	CH ₂	O	2-Cl	H	H	Me	178-180
124	2-indolyl	-	O	2-Cl	H	H	Me	> 260
125	3-indolyl	CH ₂	O	2-Cl	H	H	Me	190-192
126	4-HO-Ph	C ₂ H ₄	O	2-Cl	H	H	Me	175-180
127	2-F ₃ C-Ph	-	O	2-Cl	H	H	Me	180-182
128	4-HO-Ph	CH ₂	O	2-Cl	H	H	Me	188-190
129	5-MeO-2-indolyl	-	O	2-Cl	H	H	Me	> 250
130	Ph	-	O	2-Cl	H	H	Me	246-248
131	Ph	-	S	2-Cl	H	H	Me	200-202

132	5-MeO-2-indolyl	-	S	2-Cl	H	H	Me	> 260
133	2-NO ₂ -Ph	CH ₂	O	2-Cl	H	H	Me	204-206
134	2-F ₃ C-Ph	CH ₂	S	2-Cl	H	H	Me	> 260
135	2-NO ₂ -4-MeO-Ph	NH	S	4-Cl	H	H	Me	185-190
136	2-NO ₂ -Ph	CH ₂	S	2-Cl	H	H	Me	180-185
137	2-NO ₂ -4-MeO-Ph	NH	S	2-MeO	H	H	Bu	220-224
138	2-NO ₂ -4-MeO-Ph	NH	S	2-MeO	H	H	Bz	150-154
139	2-NO ₂ -4-MeO-Ph	NH	S	2-Me	H	H	Bu	191-193
140	2-NO ₂ -4-MeO-Ph	NH	S	2-Me	H	H	Bz	175
141	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	Ph-Ph	237-240
142	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	cyclohexyl methyl	204-206
143	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	(Me) ₂ N-C ₂ H ₄	179-182
144	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	3-HO-Bz	212-217
145	2-pyridyl	NH	S	2-Cl	H	H	Me	216-218
146	Ph	S	S	2-Cl	H	H	Me	264-266
147	Ph	O	O	2-Cl	H	H	Me	178-180
148	2-NO ₂ -4-MeO-Ph	NH	S	2-Cl	H	H	heptyl	198-200

Pharmacological study

Study of the bond to somatostatin receptors

The affinity of the compounds of the invention on human somatostatin receptors is determined by measurement of the inhibition of the bond of somatostatin-14 labelled with iodine-125 ([125I-Tyr11]SRIF-14) on the receptors of transfected CHO-K11 cells.

The human genes coding for each of the sub-types of somatostatin receptors, sst1, sst2, sst3, sst4 and sst5, have been isolated and sub-cloned (*Proc. Natl. Acad. Sci. USA* 1992, 89, 251-255; *J. Biol. Chem.* 1992, 267, 20422-20428; *Mol. Pharmacol.* 1992, 42, 2136-2142; *Proc. Natl. Acad. Sci. USA* 1993, 90, 4196-4200; *Mol. Pharmacol.* 1994, 46, 291-298). The expression vectors were constructed and the cloned cell lines were obtained by transfection in mammalian CHO-K1 cells. The plasmid pRSV-neo was included as a selection factor.

The CHO-K1 cells which express in a stable fashion the human somatostatin receptors are cultured in an RPMI 1640 medium containing 10% of foetal calf serum and 0.4 mg/ml of geneticin. The cells are collected with EDTA at 0.5 mM and centrifuged at 500 g for 5 minutes at 4°C. The pellet is resuspended in Tris 50 mM, pH 7.4 and centrifuged twice at 500 g for 5 minutes at 4°C. The cells are lysed by sonication then centrifuged at 39000 g for 10 minutes at 4°C. The pellet is resuspended in the same buffer and centrifuged at 50000 g for 5 minutes at 4°C. The cell membranes obtained are stored at -80°C until the day of the experiments.


The competitive inhibition experiments of the bond of [125I-Tyr11]SRIF-14 are carried out in duplicate in 96-well plates. The cell membranes at 10 (sst2 and sst5) or 20 (sst1, sst3 and sst4) µg of proteins/well, were incubated with [125I-Tyr11]SRIF-14 at 0.05 nM (sst2) or 0.1 nM (sst1, sst3, sst4 or sst5) for 500 (sst3), 60 (sst1 and sst2), 70 (sst5) or 90 (sst4) minutes at 37°C in a HEPES 50 mM, pH 7.4, BSA 0.2 %, MgCl₂ 5 mM, Trasylol 200 KIU/ml, bacitracin 0.02 mg/ml, phenylmethylsulphonyl fluoride 0.02 mg/ml buffer.

After the incubation period, [125I-Tyr11]SRIF-14 free or bound to the somatostatin receptors is separated on a filtration unit (Filtermate 196, Packard) with Unifilter GF/C (Packard) filter plates pretreated with 0.1 % polyethylenimine. After washing with HEPES 50 mM, the radioactivity present on the filters is measured using a Top Count (Packard) counter.

The specific bond is obtained by subtracting the non-specific bond (determined in the presence of 0.1 µM of somatostatin-14) from the total bond. The results are analyzed by non-linear regression (MDL) and the inhibition constants (K_i) determined are comprised between 10 and 10000 nM.

CLAIMS

1. Use of a compound of general formula I



in which

5 W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

R represents an aryl or heteroaryl radical, the aryl and heteroaryl radicals being optionally substituted;

X represents a radical of formula $-(CH_2)_n-Z$ in which Z represents a covalent bond, NH, O or S and n an integer of 0 to 2;

10 Y represents O or S;

R₁ represents one or more groups, identical or different, chosen from: the hydrogen atom, the hydroxy, halo radical, a lower alkyl, lower alkoxy radical;

15 R_{2a} and R_{2b} represent, independently, the hydrogen atom; a lower alkyl, lower alkenyl or lower alkynyl radical, the alkyl, alkenyl and alkynyl radicals being optionally substituted; or an R₂₁Z₂₁- radical in which Z₂₁ represents O, C(O), OC(O), S, and R₂₁ represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical;

20 R₃ represents the hydrogen atom, the halo, nitro or cyano radical, an alkyl radical with 1 to 10 carbon atoms, lower alkenyl, lower alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, lower aryloxyalkyl, heteroaryl or heteroarylalkyl, the alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted; or an R₃₁Z₃₁-

radical in which Z₃₁ represents O, C(O), OC(O), S, and R₃₁ represents the hydrogen atom, a lower alkyl, aryl or lower arylalkyl radical;

or a salt of this product, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved.

5 2. Use of a compound according to claim 1, characterized in that

W represents the hydrogen atom or a radical of formula R-X-C(Y)-;

10 R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals: lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, aryl, aryloxy, cycloalkyl or heterocycloalkyl;

15 R₁ represents one or more identical or different groups, chosen from: the hydrogen atom, the hydroxy radical, halo, a lower alkyl radical, lower alkoxy, the alkyl and alkoxy radicals being optionally substituted by one or more identical or different radicals chosen from the following radicals: trifluoromethyl, lower alkoxy, amino, lower alkyl amino and lower dialkyl amino;

R_{2a} and R_{2b} represent, independently:

the hydrogen atom;

20 a lower alkyl, lower alkenyl or lower alkynyl radical, the alkyl, alkenyl and alkynyl radicals being optionally substituted by one or more identical or different radicals, chosen from:

25 halo; an -NR₂₂R₂₃ radical in which R₂₂ and R₂₃ represent, independently, the hydrogen atom, a lower alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl, arylsulphonyl, lower alkoxy carbonyl, aryloxy carbonyl, alkylcarbonyl, arylcarbonyl or cycloalkylcarbonyl radical; or a -Z₂₂R₂₄ radical in which
30 Z₂₂ represents O,S, C(O), OC(O) and R₂₄ represents a hydrogen atom, a lower alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl or arylsulphonyl radical;

an $R_{21}Z_{21}$ - radical in which Z_{21} represents O, C(O), OC(O), S, and R_{21} represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical;

R_3 represents:

the hydrogen atom, the halo, nitro or cyano radical;

5 an alkyl radical with 1 to 10 carbon atoms, lower alkenyl, lower alkynyl, cycloalkyl, cycloalkylalkyl, aryl, lower arylalkyl, lower aryloxyalkyl, heteroaryl or lower heteroarylalkyl, the alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted by one or more identical or different radicals chosen from:

10 halo; aryl; $-NR_{32}R_{33}$ in which either R_{32} and R_{33} represent, independently, the hydrogen atom, a lower alkyl, arylalkyl or alkylcarbonyl radical, or R_{32} and R_{33} form, with the nitrogen atom to which they are attached, a heterocycloalkyl; or $-Z_{32}-R_{34}$ in which Z_{32} represents O, C(O), OC(O), S, S(O) or SO₂ and R_{34} represents the hydrogen atom, a lower alkyl, aryl or
15 lower arylalkyl radical;

a $-Z_{31}R_{31}$ radical in which Z_{31} represents O, C(O), OC(O), S, and R_{31} represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical.

3. Use of a compound according to one of claims 1 to 2, characterized in that

W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

20 R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals: lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethyloxy, hydroxy, nitro, cyano, aryl, aryloxy or heterocycloalkyl;

25 R_1 represents one or more identical or different groups, chosen from: the hydrogen atom, a halo, lower alkyl or lower alkoxy radical;

R_{2a} and R_{2b} represent, independently the hydrogen atom or a lower alkyl radical;

R_3 represents the hydrogen atom; an alkyl radical with 1 to 10 carbon atoms, cycloalkylalkyl, aryl, lower arylalkyl or lower heteroarylalkyl, the alkyl, cycloalkyl, aryl
30 and heteroaryl radicals being optionally substituted by one or more identical or different radicals chosen from:

aryl; -NR₃₂R₃₃ in which either R₃₂ and R₃₃ represent, independently, the hydrogen atom or a lower alkyl radical; or -Z₃₂-R₃₄ in which Z₃₂ represents O and R₃₄ represents the hydrogen atom or a lower alkyl radical.

4. Use of a compound according to one of the claims 1 to 3, characterized in that

5 W represents the hydrogen atom or a radical of formula R-X-C(Y)-;

R represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the following radicals: methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino;

X represents CH₂, C₂H₄, CH₂NH, NH, O, S or a covalent bond;

Y represents O or S;

15 R₁ represents one or more identical or different groups, chosen from: the hydrogen atom, a chloro, methyl or methoxy radical;

R_{2a} and R_{2b} represent, independently, the hydrogen atom or a methyl radical;

20 R₃ represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy, phenethyl, naphthylmethyl or indolylmethyl radical.

5. Use of a compound according to one of the claims 1 to 4, characterized in that the product of formula I corresponds to one of the following formulae:

- 1-butyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

25 - 1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-methyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo

[4,3-a] [1,4] diazepine;

- 1-ethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

5 - 1-propyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-phenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-pentyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

10 - 1-hexyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-(4-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

15 - 1-(4-methoxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-(1-naphthyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-(3-indolyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

20 - 1-phenethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-diphenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

25 - 1-ethoxyethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-cyclohexylmethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-(3-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 1-(dimethylaminoethyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-methyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 5 - 1-benzyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-benzyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-methyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 10 - 1-benzyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-methyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 15 - 1-butyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-benzyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-butyl-6-(2-methoxyphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 20 - 1-heptyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 1-hexyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 25 - 1-pentyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-oxoethyl]-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-

1-thioxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine;

- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide;

5 - 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxamide;

- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylbenzyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide;

- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-benzyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxamide;

10 - phenyl ester of 6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxylic acid;

- 6-(2-chlorophenyl)-7,10-dihydro-1,4-dimethyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide;

15 - 1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide;

or its substituents R,X,Y, R₁, R_{2a}, R_{2b} and R₃ are respectively the following:

- 2-MeO-Ph; NH; S; 2-Cl; H; H; Me;

- 2-Me-Ph; NH; S; 2-Cl; H; H; Me;

- 2-isoPr-Ph; NH; S; 2-Cl; H; H; Me;

20 - 2-NC-Ph; NH; S; 2-Cl; H; H; Me;

- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Et;

- 2-F₃C-Ph; NH; S; 2-Cl; H; H; H;

- 2-terBu-Ph; NH; S; 2-Cl; H; H; Me;

- 1-naphthyl; NH; S; 2-Cl; H; H; Me;

25 - 2-F₃CO-Ph; NH; S; 2-Cl; H; H; Me;

- 2-Cl-Ph; NH; S; 2-Cl; H; H; Me;

- 2-F-Ph; NH; S; 2-Cl; H; H; Me;

- 2-Et-Ph; NH; S; 2-Cl; H; H; Me;

- 2-PhO-Ph; NH; S; 2-Cl; H; H; Me;

30 - 2-Pr-Ph; NH; S; 2-Cl; H; H; Me;


- 2-Br-Ph; NH; S; 2-Cl; H; H; Me;

- 2-EtOC(O)-Ph; NH; S; 2-Cl; H; H; Me;

- 2-MeS-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,4-(MeO)-Ph; NH; S; 2-Cl; H; H; Me;
5 - 2-Cl-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Me-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,3-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2,5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
10 - 2-Me-5-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-MeO-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
15 - 2,5-Br-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Cl-5-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Pr;
- 2-F₃C-Ph; NH; S; 2-Cl; H; H; Bu;
- 2-F₃C-Ph; NH; S; H; H; H; Me;
20 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Pr;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Bu;
- 2-MeSO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
25 - 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Bz;
- 2-F₃C-Ph; NH; S; 4-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; pentyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; hexyl;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Bz;
30 - 2-NO₂-4-F-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 1-naphthyl-methyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-indolyl-methyl;
- 2-MeS-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
35 - 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Me-Ph; NH; S; 2-Cl; H; H; Me;

- 2-NO₂-4-EtO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-MeO-Bz;
- 2-NO₂-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 5 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-HO-Bz;
- 2-F₃C-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; H; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-C₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; EtOC₂H₄;
- 10 - 3-NO₂-2-pyridinyl; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; -, O; 2-Cl; H; H; Me;
- Ph; -, S; 2-Cl; H; H; Me;
- 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Me;
- 15 - 2-NO₂-Ph; CH₂; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bu;
- 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bu;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bz;
- 20 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-Ph;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; cyclohexylmethyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; (Me)₂NC₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-HO-Bz;
- Ph; S; S; 2-Cl; H; H; Me;
- 25 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; heptyl;

6. Compound of general formula II in which



in which

W' represents the hydrogen atom or a radical of formula $R'-X'-C(Y')-$;

R' represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the following radicals: methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethyloxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino;

X' represents CH_2 , C_2H_4 , CH_2NH , NH, O, S or a covalent bond;

Y' represents O or S;

R'₁ represents one or more identical or different groups, chosen from: the hydrogen atom, a chloro, methyl or methoxy radical;

R_{2a}' and R_{2b}' represent, independently, the hydrogen atom, a methyl radical;

R'₃ represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy, phenethyl, naphthylmethyl or indolylmethyl radical.

with the exclusion of the compounds of formula II in which

W' represents the hydrogen atom, R'₁ represents the chloro radical in ortho position; R_{2a}' represents the hydrogen atom and R_{2b}' represents the hydrogen atom or the methyl radical; R'₃ represents the radical methyl; and

W' represents a radical of formula $R'-X'-C(Y')-$ and

- X' represents NH; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a}' and R_{2b}' represent the hydrogen atom; R'₃ represents the methyl radical; R' represents the 4-terbutylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 2,3-dichlorophenyl, 2,4-(difluoro)phenyl, 4-phenoxy-phenyl, pyridinyl, cyanophenyl;

- X' represents NH; Y' represents S; R'₁ represents the chloro radical in ortho position; R_{2a}' and R_{2b}' represent the hydrogen atom; R'₃ represents the methyl radical; R' represents the 4-terbutylphenyl, 2,4-diterbutylphenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 4-fluorophenyl, 4-(methylsulphonyl)phenyl;

- X' represents CH₂NH; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represents the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl;

5 - X' represents the oxygen atom or a covalent bond; Y' represents O; R₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents pyridyl or cyanophenyl;

- X' represents CH₂; Y' represents O; R₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl or 4-fluorophenyl;

10 - X' represents C₂H₄; Y' represents O; R₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represent the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl.

7. Compound of general formula II according to claim 6, in which W' represents R'-X'-C(Y')- and the substituents R', X', Y', R'₁, R_{2a'}, R_{2b'} and R'₃ represent respectively:

15 - 2-F₃C-Ph; CH₂; O; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; NH; O; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; CH₂NH; S; 2-Cl; H; H; Me;

- Ph; O; O; 2-Cl; H; H; Me;
 20 - 2-F₃C-Ph; NH; S; 2-Cl; Me; H; Me;
 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; Bz;
 - 3-F₃C-Ph; NH; O; 2-Cl; H; H; Me;
 - 4-F₃C-Ph; NH; O; 2-Cl; H; H; Me;
 - 2-isoPr-Ph; NH; S; 2-Cl; H; H; Me;

25 - 2-NC-Ph; NH; S; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; Et;
 - 2-F₃C-Ph; NH; S; 2-Cl; H; H; H;
 - 2-terBu-Ph; NH; S; 2-Cl; H; H; Me;
 - 1-naphthyl; NH; S; 2-Cl; H; H; Me;
 30 - 2-Ph-Ph; NH; S; 2-Cl; H; H; Me;
 - 2-F₃CO-Ph; NH; S; 2-Cl; H; H; Me;

- 2-Cl-Ph; NH; S; 2-Cl; H; H; Me;
 - 2-F-Ph; NH; S; 2-Cl; H; H; Me;
 - 2-Et-Ph; NH; S; 2-Cl; H; H; Me;
 35 - 2-PhO-Ph; NH; S; 2-Cl; H; H; Me;

- 2-MeSO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 2-F₃C-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Bz;
- 2-F₃C-Ph; NH; S; 4-Cl; H; H; Me;
- 5 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; pentyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; hexyl;
- 3,5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Bz;
- 2-NO₂-4-F-Ph; NH; S; 2-Cl; H; H; Me;
- 10 - 2-NO₂-4-NC-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 1-naphthyl-methyl;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-indolyl-methyl;
- 2-MeS-5-F₃C-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 3-Cl; H; H; Me;
- 15 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-HO-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-5-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-EtO-Ph; NH; S; 2-Cl; H; H; Me;
- 20 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-MeO-Bz;
- 2-NO₂-4-Cl-Ph; NH; S; 2-Cl; H; H; Me;
- 2-Br-4-Me-Ph; NH; S; 2-Cl; H; H; Me;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 4-HO-Bz;
- 2-F₃C-4-NO₂-Ph; NH; S; 2-Cl; H; H; Me;
- 25 - 2-NO₂-4-MeO-Ph; NH; S; H; H; H; Bz;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-C₂H₄;
- 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; EtOC₂H₄;
- 3-NO₂-2-pyridyl; NH; S; 2-Cl; H; H; Me;
- 4-MeO-Ph; CH₂; O; 2-Cl; H; H; Me;
- 30 - 2-indolyl; -; O; 2-Cl; H; H; Me;
- 3-indolyl; CH₂; O; 2-Cl; H; H; Me;
- 4-HO-Ph; C₂H₄; O; 2-Cl; H; H; Me;
- 2-F₃C-Ph; -; O; 2-Cl; H; H; Me;
- 4-HO-Ph; CH₂; O; 2-Cl; H; H; Me;
- 35 - 5-MeO-2-indolyl; -; O; 2-Cl; H; H; Me;
- Ph; -; O; 2-Cl; H; H; Me;
- Ph; -; S; 2-Cl; H; H; Me;
- 5-MeO-2-indolyl; -; S; 2-Cl; H; H; Me;

- 2-NO₂-Ph; CH₂; O; 2-Cl; H; H; Me;
 - 2-F₃C-Ph; CH₂; S; 2-Cl; H; H; Me;
 - 2-NO₂-4-MeO-Ph; NH; S; 4-Cl; H; H; Me;
 - 2-NO₂-Ph; CH₂; S; 2-Cl; H; H; Me;
 - 5 - 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bu;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-MeO; H; H; Bz;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bu;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Me; H; H; Bz;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; Ph-Ph;
 - 10 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; cyclohexyl methyl;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; (Me)₂NC₂H₄;
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; 3-HO-Bz;
 - 2-pyridyl; NH; S; 2-Cl; H; H; Me;
 - Ph; S; S; 2-Cl; H; H; Me;
 - 15 - Ph; O; S; 2-Cl; H; H; Me,
 - 2-NO₂-4-MeO-Ph; NH; S; 2-Cl; H; H; heptyl,
- but also the compounds of formula II in which W' represents the hydrogen atom and the substituents R'₁, R'_{2a}, R'_{2b} and R'₃ represent respectively:
- 2-Cl; H; H; butyl;
 - 20 - 2-Cl; H; H; benzyl;
 - 2-Cl; H; H; H;
 - 2-Cl; H; H; ethyl;
 - 2-Cl; H; H; propyl;
 - 2-Cl; H; H; Ph;
 - 25 - 2-Cl; H; H; pentyl;
 - 2-Cl; H; H; hexyl;
 - 2-Cl; H; H; 4-HO-Bz;
 - 2-Cl; H; H; 4-MeO-Bz;
 - 2-Cl; H; H; 1-naphthyl-methyl;
 - 30 - 2-Cl; H; H; 3-indolyl-methyl;
 - 2-Cl; H; H; Ph-C₂H₄;
 - 2-Cl; H; H; Ph-Ph;
 - 2-Cl; H; H; EtOC₂H₄;
 - 2-Cl; H; H; cyclohexylmethyl;
 - 35 - 2-Cl; H; H; 3-OH-Bz;
 - 2-Cl; H; H; (Me)₂NC₂H₄;
 - H; H; H; Me;
 - 4-Cl; H; H; Bz;


- H; H; H; Bz;
- 4-Cl; H; H; Me;
- 3-Cl; H; H; benzyl;
- 3-Cl; H; H; Me;
- 5 - 2-Me; H; H; butyl;
- 2-Me; H; H; benzyl;
- 2-MeO; H; H; butyl;
- 2-Cl; H; H; heptyl;
- 4-Cl; H; H; hexyl;
- 10 - 4-Cl; H; H; pentyl;

8. As a medicament, a compound of general formula II according to one of claims 6 to 7.

9. Therapeutic composition containing, as active ingredient, at least one medicament according to claim 8, in combination with a pharmaceutically acceptable support.

CLAIMS

1. Use of a compound of general formula I



in which

5 W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals : lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethyloxy, hydroxy, nitro, cyano, aryl, aryloxy, cycloalkyl or heterocycloalkyl ;

X represents a radical of formula $-(CH_2)_n-Z$ in which Z represents a covalent bond, NH, O or S and n an integer of 0 to 2 ;

Y represents O or S ;

15 R₁ represents one or more groups, identical or different, chosen from : the hydrogen atom, the hydroxy, halo radical, a lower alkyl, lower alkoxy radical, the alkyl and alkoxy radicals being optionally substituted by one or more identical or different radicals chosen from the following radicals : trifluoromethyl, lower alkoxy, amino, lower alkyl amino and lower dialkyl amino ;

R_{2a} and R_{2b} represent, independently :

20 the hydrogen atom ;

a lower alkyl, lower alkenyl or lower alkynyl radical, the alkyl, alkenyl and alkynyl radicals being optionally substituted by one or more identical or different radicals, chosen from :

- 5 halo ; an $-NR_{22}R_{23}$ radical in which R_{22} and R_{23} represent, independently, the hydrogen atom, a lower alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl, arylsulphonyl, lower alkoxy carbonyl, aryloxy carbonyl, alkylcarbonyl, arylcarbonyl or cycloalkylcarbonyl radical; or a $-Z_{22}R_{24}$ radical in which Z_{22} represents O, S, C(O), OC(O) and R_{24} represents a hydrogen atom, a lower
10 alkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, alkylsulphonyl, cycloalkylsulphonyl or arylsulphonyl radical ;

an $R_{21}Z_{21}$ - radical in which Z_{21} represents O, C(O), OC(O), S, and R_{21} represents the hydrogen atom, a lower alkyl, aryl or arylalkyl radical ;

R_3 represents :

- 15 the hydrogen atom, the halo, nitro or cyano radical ;
- an alkyl radical with 1 to 10 carbon atoms, lower alkenyl, lower alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, lower aryloxyalkyl, heteroaryl or heteroarylalkyl, the alkyl, alkenyl, alkynyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted by one or more identical or different radicals chosen
20 from :

- 25 halo ; aryl ; $-NR_{32}R_{33}$ in which either R_{32} and R_{33} represent, independently, the hydrogen atom, a lower alkyl, arylalkyl or alkylcarbonyl radical, or R_{32} and R_{33} form, with the nitrogen atom to which they are attached, a heterocycloalkyl ; or $-Z_{32}-R_{34}$ in which Z_{32} represents O, C(O), OC(O), S, S(O) or SO₂ and R_{34} represents the hydrogen atom, a lower alkyl, aryl or lower arylalkyl radical ;

an $R_{31}Z_{31}$ - radical in which Z_{31} represents O, C(O), OC(O), S, and R_{31} represents the hydrogen atom, a lower alkyl, aryl or lower arylalkyl radical ;

- 30 or a salt of this product, for the preparation of a medicament intended to treat the pathological states or the diseases in which one (or more) of the somatostatin receptors is involved,

it being understood that the term "lower", when it refers to an alkyl, alkoxy, alkylthio, alkenyl or alkynyl radical, means that the radical contains up to 6 carbon atoms.

2. Use of a compound according to claim 1, characterized in that

W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

5 R represents an aryl or heteroaryl radical, the aryl or heteroaryl radical being optionally substituted by one or more identical or different substituents, chosen from the following radicals : lower alkyl, lower alkoxy, lower alkylthio, lower alkoxy carbonyl, lower alkyl sulphonyl, halo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, aryl, aryloxy or heterocycloalkyl ;

10 R_1 represents one or more identical or different groups, chosen from : the hydrogen atom, a halo, lower alkyl or lower alkoxy radical ;

R_{2a} and R_{2b} represent, independently, the hydrogen atom or a lower alkyl radical ;

15 R_3 represents the hydrogen atom ; an alkyl radical with 1 to 10 carbon atoms, cycloalkylalkyl, aryl, lower arylalkyl or heteroarylalkyl, the alkyl, cycloalkyl, aryl and heteroaryl radicals being optionally substituted by one or more identical or different radicals, chosen from :

aryl ; $-NR_{32}R_{33}$ in which either R_{32} and R_{33} represent, independently, the hydrogen atom or a lower alkyl radical ; or $-Z_{32}-R_{34}$ in which Z_{32} represents O and R_{34} represents the hydrogen atom or a lower alkyl radical.

20 3. Use of a compound according to one of the claims 1 to 2, characterized in that

W represents the hydrogen atom or a radical of formula $R-X-C(Y)-$;

25 R represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the following radicals : methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino ;

X represents CH_2 , C_2H_4 , CH_2NH , NH , O, S or a covalent bond ;

Y represents O or S ;

R₁ represents one or more identical or different groups, chosen from : the hydrogen atom, a chloro, methyl or methoxy radical ;

R_{2a} and R_{2b} represent, independently, the hydrogen atom or a methyl radical ;

5 R₃ represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy, phenethyl, naphthylmethyl or indolylmethyl radical.

4. Use of a compound according to one of the claims 1 to 3, characterized in that the product of formula I corresponds to one of the following formulae

- 10 - 1-butyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-benzyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 15 - 1-methyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-ethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 20 - 1-propyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-phenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 25 - 1-pentyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-hexyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-(4-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;

- 1-(4-methoxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-(1-naphthyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 5 - 1-(3-indolyl-methyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-phenethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 10 - 1-diphenyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-ethoxyethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-cyclohexylmethyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 15 - 1-(3-hydroxybenzyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-(dimethylaminoethyl)-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 20 - 1-methyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-benzyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-benzyl-6-phenyl-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 25 - 1-methyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-benzyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;

- 1-methyl-6-(3-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-butyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 5 - 1-benzyl-6-(2-methylphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-butyl-6-(2-methoxyphenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 10 - 1-heptyl-6-(2-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-hexyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 1-pentyl-6-(4-chlorophenyl)-7,8,9,10-tetrahydro-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 15 - 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-oxoethyl]-4H-pyrido[4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 6-(2-chlorophenyl)-7,8,9,10-tetrahydro-1-methyl-9-[2-(2-trifluoromethylphenyl)-1-thioxoethyl]-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine ;
- 20 - 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide ;
- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carboxamide ;
- 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-(2-trifluoromethylbenzyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine-9(8H)-carbothioamide ;
- 25 - 6-(2-chlorophenyl)-7,10-dihydro-1-methyl-N-benzyl-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxamide ;
- phenyl ester of 6-(2-chlorophenyl)-7,10-dihydro-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carboxylic acid ;

MODIFIED SHEET

- 6-(2-chlorophenyl)-7,10-dihydro-1,4-dimethyl-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide ;

- 1-benzyl-6-(2-chlorophenyl)-7,10-dihydro-N-(2-trifluoromethylphenyl)-4H-pyrido [4',3';4,5] thieno [3,2-f] [1,2,4] triazolo [4,3-a] [1,4] diazepine -9(8H)-carbothioamide ;

5 or its substituents R,X,Y, R₁, R_{2a}, R_{2b} and R₃ are respectively the following :

- 2-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-isoPr-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-NC-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

10 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Et ;

- 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; H ;

- 2-terBu-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 1-naphthyl ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-F₃CO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

15 - 2-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Et-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-PhO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Pr-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

20 - 2-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-EtOC(O)-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-MeS-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-MeO-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

25 - 2,4-(MeO)-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Cl-5-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Me-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2,3-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2,5-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

30 - 2,5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-Me-5-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-F₃C-4-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-NO₂-4-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

35 - 2-MeO-4-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2,5-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;


MODIFIED SHEET

009901422 30-05-2000

- 2-Cl-5-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Pr ;
- 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Bu ;
- 2-F₃C-Ph ; NH ; S ; H ; H ; H ; Me ;
- 5 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Pr ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Bu ;
- 2-MeSO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-4-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 10 - 2-NO₂-4-MeO-Ph ; NH ; S ; 4-Cl ; H ; H ; Bz ;
- 2-F₃C-Ph ; NH ; S ; 4-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; pentyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; hexyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 3-Cl ; H ; H ; Bz ;
- 15 - 2-NO₂-4-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-NC-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 1-naphthyl-methyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 3-indolyl-methyl ;
- 2-MeS-5-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 20 - 2-NO₂-4-MeO-Ph ; NH ; S ; 3-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-5-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-EtO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 25 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 4-MeO-Bz ;
- 2-NO₂-4-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Br-4-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 4-HO-Bz ;
- 2-F₃C-4-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 30 - 2-NO₂-4-MeO-Ph ; NH ; S ; H ; H ; H ; Bz ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph-C₂H₄ ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; EtOC₂H₄ ;
- 3-NO₂-2-pyridinyl ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; - ; O ; 2-Cl ; H ; H ; Me ;
- 35 - Ph ; - ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; CH₂ ; S ; 2-Cl ; H ; H ; Me ;

- 2-NO₂-4-MeO-Ph ; NH ; S ; 4-Cl ; H ; H ; Me ;
- 2-NO₂-Ph ; CH₂ ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-MeO ; H ; H ; Bu ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-MeO ; H ; H ; Bz ;
- 5 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Me ; H ; H ; Bu ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Me ; H ; H ; Bz ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph-Ph ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; cyclohexylmethyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; (Me)₂NC₂H₄ ;
- 10 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 3-HO-Bz ;
- Ph ; S ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; heptyl.

5. Compound of general formula II in which



15 in which

W' represents the hydrogen atom or a radical of formula R'-X'-C(Y')- ;

R' represents the phenyl, naphthyl, indolyl or pyridyl radical, these radicals being optionally substituted by one or more identical or different substituents chosen from the following radicals : methyl, ethyl, propyl, isopropyl, butyl, ter-butyl, methoxy, ethoxy, methylthio, ethylthio, methoxycarbonyl, ethoxycarbonyl, methylsulphonyl, ethylsulphonyl, chloro, fluoro, bromo, trifluoromethyl, trifluoromethoxy, hydroxy, nitro, cyano, phenyl, phenoxy or morpholino ;

X' represents CH₂, C₂H₄, CH₂NH, NH, O, S or a covalent bond ;

Y' represents O or S ;

R'₁ represents one or more identical or different groups, chosen from : the hydrogen atom, a chloro, methyl or methoxy radical ;

R_{2a'} and R_{2b'} represent, independently, the hydrogen atom, a methyl radical ;

5 R'₃ represents the hydrogen atom, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, methoxyethyl, ethoxyethyl, dimethylaminoethyl, cyclohexylmethyl, phenyl, diphenyl, benzyl optionally substituted by the hydroxy or methoxy, phenethyl, naphthylmethyl or indolylmethyl radical.

with the exclusion of the compounds of formula II in which

10 W' represents the hydrogen atom, R'₁ represents the chloro radical in ortho position ; R_{2a'} represents the hydrogen atom and R_{2b'} represents the hydrogen atom or the methyl radical ; R'₃ represents the radical methyl ; and

W' represents a radical of formula R'-X'-C(Y')- and

15 - X' represents NH ; Y' represents O ; R'₁ represents the chloro radical in ortho position ; R_{2a'} and R_{2b'} represent the hydrogen atom ; R'₃ represents the methyl radical ; R' represents the 4-terbutylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 2,3-dichlorophenyl, 2,4-(difluoro)phenyl, 4-phenoxy-phenyl, pyridinyl, cyanophenyl ;

20 - X' represents NH ; Y' represents S ; R'₁ represents the chloro radical in ortho position ; R_{2a'} and R_{2b'} represent the hydrogen atom ; R'₃ represents the methyl radical ; R' represents the 4-terbutylphenyl, 2,4-diterbutylphenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4,5-trimethoxyphenyl, 4-fluorophenyl, 4-(methylsulphonyl)phenyl ;

25 - X' represents CH₂NH; Y' represents O; R'₁ represents the chloro radical in ortho position; R_{2a'} and R_{2b'} represents the hydrogen atom; R'₃ represents the methyl radical; R' represents phenyl;

30 - X' represents the oxygen atom or a covalent bond ; Y' represents O ; R'₁ represents the chloro radical in ortho position ; R_{2a'} and R_{2b'} represent the hydrogen atom ; R'₃ represents the methyl radical ; R' represents pyridyl or cyanophenyl ;

- X' represents CH₂ ; Y' represents O ; R₁ represents the chloro radical in ortho position ; R_{2a'} and R_{2b'} represent the hydrogen atom ; R'₃ represents the methyl radical ; R' represents phenyl or 4-fluorophenyl ;

- 5 - X' represents C₂H₄ ; Y' represents O ; R₁ represents the chloro radical in ortho position ; R_{2a'} and R_{2b'} represent the hydrogen atom ; R'₃ represents the methyl radical ; R' represents phenyl.

6. Compound of general formula II according to claim 5, in which W' represents R'-X'-C(Y')- and the substituents R', X', Y', R'₁, R_{2a'}, R_{2b'} and R'₃ represent respectively :

- 10 - 2-F₃C-Ph ; CH₂ ; O ; 2-Cl ; H ; H ; Me ;
 - 2-F₃C-Ph ; CH₂ ; S ; 2-Cl ; H ; H ; Me ;
 - 2-F₃C-Ph ; NH ; O ; 2-Cl ; H ; H ; Me ;
 - 2-F₃C-Ph ; CH₂NH ; S ; 2-Cl ; H ; H ; Me ;
 - Ph ; O ; O ; 2-Cl ; H ; H ; Me ;
 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; Me ; H ; Me ;
 15 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Bz ;
 - 3-F₃C-Ph ; NH ; O ; 2-Cl ; H ; H ; Me ;
 - 4-F₃C-Ph ; NH ; O ; 2-Cl ; H ; H ; Me ;
 - 2-isoPr-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-NC-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 20 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Et ;
 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; H ;
 - 2-terBu-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 1-naphthyl ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-Ph-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 25 - 2-F₃CO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-Et-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-PhO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 30 - 2-Pr-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-EtO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-EtOC(O)-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 35 - 2-MeS-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
 - 2-morpholino-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;

- 2-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,6-isoPr-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,6-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,5-(MeO)-Ph ; NH ; O ; 2-Cl ; H ; H ; Me ;
- 5 - 2-MeO-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,4-(MeO)-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Cl-5-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Me-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,3-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 10 - 2,5-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2,5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Cl-4-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Me-3-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Me-5-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 15 - 2,3-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-4-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-MeO-4-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 20 - 2,5-Br-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-MeO-5-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Cl-4-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Cl-5-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Pr ;
- 25 - 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Bu ;
- 3-Ph-6-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; NH ; S ; H ; H ; H ; Me ;
- 2-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Pr ;
- 30 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Bu ;
- 2-NO₂-4-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-MeSO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-4-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 4-Cl ; H ; H ; Bz ;
- 35 - 2-F₃C-Ph ; NH ; S ; 4-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; pentyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; hexyl ;

- 3,5-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 3-Cl ; H ; H ; Bz ;
- 2-NO₂-4-F-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-NC-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 5 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 1-naphthyl-methyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 3-indolyl-methyl ;
- 2-MeS-5-F₃C-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 3-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 10 - 2-NO₂-4-HO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-5-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-5-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-EtO-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 4-MeO-Bz ;
- 15 - 2-NO₂-4-Cl-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-Br-4-Me-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 4-HO-Bz ;
- 2-F₃C-4-NO₂-Ph ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; H ; H ; H ; Bz ;
- 20 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph-C₂H₄ ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; EtOC₂H₄ ;
- 3-NO₂-2-pyridyl ; NH ; S ; 2-Cl ; H ; H ; Me ;
- 4-MeO-Ph ; CH₂ ; O ; 2-Cl ; H ; H ; Me ;
- 2-indolyl ; - ; O ; 2-Cl ; H ; H ; Me ;
- 25 - 3-indolyl ; CH₂ ; O ; 2-Cl ; H ; H ; Me ;
- 4-HO-Ph ; C₂H₄ ; O ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; - ; O ; 2-Cl ; H ; H ; Me ;
- 4-HO-Ph ; CH₂ ; O ; 2-Cl ; H ; H ; Me ;
- 5-MeO-2-indolyl ; - ; O ; 2-Cl ; H ; H ; Me ;
- 30 - Ph ; - ; O ; 2-Cl ; H ; H ; Me ;
- Ph ; - ; S ; 2-Cl ; H ; H ; Me ;
- 5-MeO-2-indolyl ; - ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-Ph ; CH₂ ; O ; 2-Cl ; H ; H ; Me ;
- 2-F₃C-Ph ; CH₂ ; S ; 2-Cl ; H ; H ; Me ;
- 35 - 2-NO₂-4-MeO-Ph ; NH ; S ; 4-Cl ; H ; H ; Me ;
- 2-NO₂-Ph ; CH₂ ; S ; 2-Cl ; H ; H ; Me ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-MeO ; H ; H ; Bu ;

- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-MeO ; H ; H ; Bz ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Me ; H ; H ; Bu ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Me ; H ; H ; Bz ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; Ph-Ph ;
- 5 - 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; cyclohexyl methyl ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; (Me)₂NC₂H₄ ;
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; 3-HO-Bz ;
- 2-pyridyl ; NH ; S ; 2-Cl ; H ; H ; Me ;
- Ph ; S ; S ; 2-Cl ; H ; H ; Me ;
- 10 - Ph ; O ; S ; 2-Cl ; H ; H ; Me,
- 2-NO₂-4-MeO-Ph ; NH ; S ; 2-Cl ; H ; H ; heptyl,
- but also the compounds of formula II in which W' represents the hydrogen atom and the substituents R'₁, R'_{2a}, R'_{2b} and R'₃ represent respectively :
- 2-Cl ; H ; H ; butyl ;
- 15 - 2-Cl ; H ; H ; benzyl ;
- 2-Cl ; H ; H ; H ;
- 2-Cl ; H ; H ; ethyl ;
- 2-Cl ; H ; H ; propyl ;
- 2-Cl ; H ; H ; Ph ;
- 20 - 2-Cl ; H ; H ; pentyl ;
- 2-Cl ; H ; H ; hexyl ;
- 2-Cl ; H ; H ; 4-HO-Bz ;
- 2-Cl ; H ; H ; 4-MeO-Bz ;
- 2-Cl ; H ; H ; 1-naphthyl-methyl ;
- 25 - 2-Cl ; H ; H ; 3-indolyl-methyl ;
- 2-Cl ; H ; H ; Ph-C₂H₄ ;
- 2-Cl ; H ; H ; Ph-Ph ;
- 2-Cl ; H ; H ; EtOC₂H₄ ;
- 2-Cl ; H ; H ; cyclohexylmethyl ;
- 30 - 2-Cl ; H ; H ; 3-OH-Bz ;
- 2-Cl ; H ; H ; (Me)₂NC₂H₄ ;
- H ; H ; H ; Me ;
- 4-Cl ; H ; H ; Bz ;
- H ; H ; H ; Bz ;
- 35 - 4-Cl ; H ; H ; Me ;
- 3-Cl ; H ; H ; benzyl ;
- 3-Cl ; H ; H ; Me ;
- 2-Me ; H ; H ; butyl ;

- 2-Me ; H ; H ; benzyl ;
- 2-MeO ; H ; H ; butyl ;
- 2-Cl ; H ; H ; heptyl ;
- 4-Cl ; H ; H ; hexyl ;
- 5 - 4-Cl ; H ; H ; pentyl.

7. As a medicament, a compound of general formula II according to one of claims 5 to 6.

8. Therapeutic composition containing, as active ingredient, at least one medicament according to claim 7, in combination with a pharmaceutically acceptable support.

10

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (8-96)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION <input checked="" type="checkbox"/> Declaration OR Submitted with Initial Filing <input type="checkbox"/> Declaration Submitted after Initial Filing	Attorney Docket Number	
	First Named Inventor	D. BIGG
	COMPLETE IF KNOWN	
	Application Number	
	Filing Date	
	Group Art Unit	
	Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

USE OF DIAZEPINS FOR PREPARING MEDICINES FOR TREATING PATHOLOGICAL
CONDITIONS OR DISEASES INVOLVING ONE OF THE GROWTH HORMONE RELEASE
INHIBITING FACTOR RECEPTORS

(Title of the invention)

the specification of which

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY)

06/15/1999

as United States Application Number or PCT International

Application Number

PCT/FR99/01422

and was amended on (MM/DD/YYYY)

(if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code §119 (a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365 (a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
98/07509	France	06/15/1998	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto:

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto.

(Page 1 of 5)

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner of Patents and Trademarks, Washington, DC 20231.

(Janvier 1997)

Please type a plus sign (+) inside this box → +

PTO/SB/01 (8-96)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Name	Registration Number	Name	Registration Number
Charles A. Muserlian	19,683		
Jordan B. Bierman	18,629		
Donald C. Lucas	31,275		
Bierman, Muserlian and Lucas	18,818		

☐ Additional registered practitioner(s) named on a supplemental sheet attached hereto.

Direct all correspondence to:

Name	Bierman, Muserlian and Lucas		
Address			
Address	600 Third Avenue		
City	New York	State	New York
		ZIP	10016
Country	U.S.A.	Telephone	(212) 661-8000
		Fax	(212) 661-8002

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name	Dennis	Middle Initial		Family Name	BIGG	Suffix e.g. Jr.	
Inventor's Signature						Date	Nov. 17, 2000
Residence: City	GIF-SUR-YVETTE	State		Country	FRANCE	Citizenship	French
Post Office Address	12 rue des Bénédictines						
Post Office Address							
City	GIF-SUR-YVETTE	State		Zip	F-91190	Country	FRANCE

☒ Additional inventors are being named on supplemental sheet(s) attached hereto

Please type a plus sign (+) inside this box → +

PTO/SB/01 (8-96)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.



DECLARATION	ADDITIONAL INVENTOR(S) Supplemental Sheet
--------------------	--

Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Anne-Marie	Middle Initial		Family Name	LIBERATORE	Suffix e.g. Jr.	
Inventor's Signature	<i>Libertore</i>				Date	20.11.2000	
Residence: City	AUFFARGIS	State		Country	FRANCE	Citizenship	French
Post Office Address	10 rue de la Croix Picard						
Post Office Address	Saint Benoist						
City	AUFFARGIS	State		Zip	F-78610	Country	FRANCE
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Jacques	Middle Initial		Family Name	POMMIER	Suffix e.g. Jr.	
Inventor's Signature	<i>Jacques Pommier</i>				Date	20.11.2000	
Residence: City	COLOMBES	State		Country	FRANCE	Citizenship	French
Post Office Address	93/95 avenue Henri Barbusse						
Post Office Address							
City	COLOMBES	State		Zip	F-92700	Country	FRANCE
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	John	Middle Initial		Family Name	TAYLOR	Suffix e.g. Jr.	
Inventor's Signature	<i>John Taylor</i>				Date	11/14/00	
Residence: City	UPTON	State	MA	Country	U.S.A.	Citizenship	U.S.
Post Office Address	74 Fiske Mill Road						
Post Office Address							
City	UPTON	State	MA	Zip	01568	Country	U.S.A.
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name		Middle Initial		Family Name		Suffix e.g. Jr.	
Inventor's Signature					Date		
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	
<input type="checkbox"/> Additional inventors are being named on supplemental sheet(s) attached hereto							

+

[illegible]

Additional provisional applications:	
Application Number	Filing Date (MM/DD/YYYY)

Additional U.S. applications:			
U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

Please type a plus sign (+) inside this box → ☐

Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

+

DECLARATION

REGISTERED PRACTITIONER INFORMATION (Supplemental Sheet)

Name	Registration Number	Name	Registration Number